arXiv:1606.06264v2 [math.RT] 16 Dec 2016

Character tables of Sylow p-subgroups ${}^3D_4^{syl}(q^3)$ of the Steinberg triality groups

Yujiao Sun

Institut für Algebra und Zahnlentheorie
Universität Stuttgart, 70569 Stuttgart, Germany

E-mail: yujiao.sun@mathematik.uni-stuttgart.de

Abstract

We determine the character tables of Sylow p-subgroups ${}^3D_4^{syl}(q^3)$ of the Steinberg triality groups ${}^3D_4(q^3)$.

1 Introduction

Let p be a fixed prime, $\mathbb{N}^*$ the set of positive integers, $q := p^k$ with a fixed $k \in \mathbb{N}^*$, $\mathbb{F}_q$ the finite field with q elements and $U_n(q)$ ($n \in \mathbb{N}^*$) the group of upper unitriangular $n \times n$-matrices (i.e. upper triangular matrices with 1s on the main diagonal) with entries in $\mathbb{F}_q$. Then $U_n(q)$ is a Sylow p-subgroup of the general linear group $GL_n(q)$ of invertible $n \times n$-matrices over $\mathbb{F}_q$.

G. Higman [Hig60] formulated a longstanding conjecture: for a fixed n, the number of conjugacy classes of $U_n(q)$ is determined by a polynomial in q with integral coefficients depending on n. G. Lehrer [Leh74] refined Higman's conjecture: the number of irreducible characters of $U_n(q)$ of degree q^c ($c \in \mathbb{N}$) is an integer polynomial in q. I. M. Isaacs [Isa95] proved that the degrees of complex irreducible characters of $\mathbb{F}_q$-algebra groups are powers of q. Then I. M. Isaacs [Isa07] gave a strengthened form of Lehrer's conjecture: the number of irreducible characters of degree q^c is some polynomial in $(q-1)$ with non-negative integer coefficients. A. Vera-López and J. M. Arregi [VLA03] proved that Higman's conjecture holds when $n \leq 13$. Recently, I. Pak and A. Soffer [PS15] proved that it holds when $n \leq 16$.

It is natural to consider Higman's conjecture, Lehrer's conjecture and Isaacs' conjecture for the Sylow p-subgroups of other finite groups of Lie type. Let $G(q)$ be a finite group of Lie type, $U(q)$ a Sylow p-subgroup of $G(q)$, $k(U(q))$ the number of conjugacy classes, $\#\mathrm{Irr}(U(q))$ the number of all complex irreducible characters, $\#\mathrm{Irr}(U(q), q^c)$ the number of complex irreducible characters of degree q^c, $\#M(U(q))$ the number of pairwise orthogonal irreducible constituents of the regular $U(q)$-module $\mathbb{C}U(q)$. S. M. Goodwin, P. Mosch and G. Röhrle [GR09, GMR14, GMR16] obtained an algorithm and calculated $k(U(q))$ for $U(q)$ of rank at most 8, except E_8.

For the Sylow p-subgroup $D_4^{syl}(q)$ of the Chevalley group $D_4(q)$ of type D_4, F. Himstedt, T. Le and K. Magaard [HLM11] determined the complex irreducible characters and calculated $\#\mathrm{Irr}(D_4^{syl}(q), q^c)$. M. Jedlitschky [Jed13, Appendix A.3] decomposed the regular module $\mathbb{C}D_4^{syl}(q)$ with $p \neq 2$ into irreducible constituents, and obtained $\#\mathrm{Irr}(D_4^{syl}(q), q^c)$ and $\#\mathrm{Irr}(D_4^{syl}(q))$. S. M. Goodwin, T. Le and K. Magaard [GLM15] constructed the *generic character table* of $D_4^{syl}(q)$. Higman's conjecture, Lehrer's conjecture and Isaacs' conjecture are true for $D_4^{syl}(q)$.

For the Sylow p-subgroup $G_2^{syl}(q)$ $(p > 3)$ of the Chevalley group $G_2(q)$ of type G_2, F. Himstedt, T. Le and K. Magaard [HLM16] determined most irreducible characters of $G_2^{syl}(q)$ by parameterizing *midafis* (minimal degree almost faithful irreducible characters).

S. M. Goodwin, T. Le, K. Magaard and A. Paolini [GLMP16] parameterized the irreducible characters of the Sylow p-subgroup $F_4^{syl}(q)$ $(p > 2)$ of the Chevalley group $F_4(q)$ of type F_4.

Let ${}^3D_4^{syl}(q^3)$ be a Sylow p-subgroup of the Steinberg triality group ${}^3D_4^{syl}(q^3)$. T. Le [Le13] constructed and counted all ordinary irreducible characters of ${}^3D_4^{syl}(q^3)$, mainly using Clifford theory. The author in [Sun16] determined a supercharacter theory for the matrix Sylow p-subgroup ${}^3D_4^{syl}(q^3)$, and established the supercharacter table of ${}^3D_4^{syl}(q^3)$.

Let p be a fixed odd prime, $q := p^k$ a fixed power of p. In this paper, we establish the conjugacy classes of ${}^3D_4^{syl}(q^3)$ (see Table 1). Based on T. Le's work [Le13], we give the specific constructions (see Proposition 4.12) of the irreducible characters of ${}^3D_4^{syl}(q^3)$ by Clifford's Theorem (see [CR81, Section 11]). Finally, we further determine the character table of ${}^3D_4^{syl}(q^3)$ (see Table 4).

Here we fix the notations: let K a field, K^* the multiplicative group $K\backslash\{0\}$ of K, K^+ the additive group of K, $\mathbb{F}_q$ the finite field with q elements, $\mathbb{F}_{q^3}$ the finite field with q^3 elements, $\mathbb{C}$ the complex field, $\mathbb{Z}$ the set of all integers, $\mathbb{N}$ the set $\{0, 1, 2, \dots\}$ of all non-negative integers, $\mathbb{N}^*$ the set $\{1, 2, \dots\}$ of all positive integers.

2 Sylow p-subgroups ${}^3D_4^{syl}(q^3)$

In this section, we recall the root system of D_4, the construction of a Sylow p-subgroup ${}^3D_4^{syl}(q^3)$ of the Steinberg triality group ${}^3D_4(q^3)$, and the commutator relations of ${}^3D_4^{syl}(q^3)$. The main references are [Car72] and [Hum78].

Let $\mathcal{V}_4$ be a fixed Euclidean space with the usual orthonormal basis ε_1, ε_2, ε_3, ε_4, where the inner product $(\ ,\ )$ is the usual one. The set $\Phi_{D_4} = \{\pm\varepsilon_i \pm \varepsilon_j \mid 1 \le i < j \le 4\}$ is a root system of type D_4, and $(r, r) = 2$ for all $r \in \Phi_{D_4}$. The fundamental system of roots of the root system Φ_{D_4} is $\Delta_{D_4} = \{\varepsilon_1 - \varepsilon_2,\ \varepsilon_2 - \varepsilon_3,\ \varepsilon_3 - \varepsilon_4,\ \varepsilon_3 + \varepsilon_4\}$. The positive system (relative to Δ_{D_4}) of roots of Φ_{D_4} is $\Phi_{D_4}^+ := \{\varepsilon_i \pm \varepsilon_j \mid 1 \le i < j \le 4\}$. Let $r_1 := \varepsilon_1 - \varepsilon_2,\ r_2 := \varepsilon_2 - \varepsilon_3,\ r_3 := \varepsilon_3 - \varepsilon_4,\ r_4 := \varepsilon_3 + \varepsilon_4$, then $\Delta_{D_4} = \{r_1, r_2, r_3, r_4\}$.

Let $r := \sum_{i=1}^4 x_i r_i \in \mathcal{V}_4$, $s := \sum_{i=1}^4 y_i r_i \in \mathcal{V}_4$, then we write $r \prec s$, if $\sum_{i=1}^4 x_i < \sum_{i=1}^4 y_i$, or if $\sum_{i=1}^4 x_i = \sum_{i=1}^4 y_i$ and the first non-zero coefficient $x_i - y_i$ is positive. If $r = \sum_{i=1}^4 x_i r_i \in \Phi_{D_4}$, the height of r (relative to Δ_{D_4}) is denoted by $\mathrm{ht}(r) := \sum_{i=1}^4 x_i$. Let $r_5 := r_1 + r_2,\ r_6 := r_2 + r_3,\ r_7 := r_2 + r_4,\ r_8 := r_1 + r_2 + r_3,\ r_9 := r_1 + r_2 + r_4,\ r_{10} := r_2 + r_3 + r_4,\ r_{11} := r_1 + r_2 + r_3 + r_4,\ r_{12} := r_1 + 2r_2 + r_3 + r_4$, then $\Phi_{D_4}^+ = \{r_i \mid i = 1, 2, \dots, 12\}$. We get the total order of $\Phi_{D_4}^+$ as follows

$$0 \prec r_1 \prec r_2 \prec r_3 \prec r_4 \prec r_5 \prec r_6 \prec r_7 \prec r_8 \prec r_9 \prec r_{10} \prec r_{11} \prec r_{12}.$$

Let $\mathcal{L}$ be a complex simple Lie algebra of type D_4 with a Cartan subalgebra $\mathcal{H}$, a Cartan decomposition $\mathcal{L} = \mathcal{H} \oplus \sum_{r \in \Phi_{D_4}} \mathcal{L}_r$, and a Chevalley basis $\{h_r \mid r \in \Delta_{D_4}\} \cup \{e_r \mid r \in \Phi_{D_4}\}$, where $e_r \in \mathcal{L}_r$ for all $r \in \Phi_{D_4}$. Then $[e_r, e_s] = N_{r,s} e_{r+s}$ for all $r, s \in \Phi_{D_4}$, where $N_{r,s} = \pm(n_{r,s} + 1)$ and $n_{r,s}$ is the greatest integer for which $s - n_{r,s} r \in \Phi_{D_4}$.

We choose the signs of the structure constants $N_{r,s}$ for extraspecial pairs $(r,\ s)$ as follows,

$$\begin{aligned} &- N_{r_1,r_2} = N_{r_2,r_3} = N_{r_2,r_4} = 1, && N_{r_1,\ r_2+r_3+r_4} = 1, \\ &N_{r_1,r_2+r_3} = N_{r_1,r_2+r_4} = N_{r_2,r_3+r_4} = 1, && N_{r_2,\ r_1+r_2+r_3+r_4} = 1. \end{aligned}$$

Then the structure constants of the other special pairs of $r_i \in \Phi_{D_4}^+ \backslash \Delta_{D_4}$ are 1.

Let $x \in \mathcal{L}$. We define a linear map by $\operatorname{ad} x : \mathcal{L} \to \mathcal{L} : y \mapsto [x, y] := xy - yx$. Let $x_r(t) := \exp(t \operatorname{ad} e_r)$ for all $r \in \Phi_{D_4}$, $t \in \mathbb{C}$. Let K be a field, then the Chevalley group of $\mathcal{L}$ over K is defined by $D_4(K) := \langle x_r(t) \mid r \in \Phi_{D_4}, t \in K \rangle$. The Chevalley group of $\mathcal{L}$ over $\mathbb{F}_q$ is also denoted by $D_4(q)$. Let $r \in \Phi_{D_4}$, then the root subgroup is denoted by $X_r := \langle x_r(t) \mid t \in K \rangle$. The subgroup $U_{D_4} := \langle x_r(t) \mid r \in \Phi_{D_4}^+, t \in \mathbb{F}_q \rangle$ is a Sylow p-subgroup of $D_4(q)$, denoted by $D_4^{syl}(q)$.

Let σ be a non-trivial symmetry of the Dynkin diagram of $\mathcal{L}_{D_4}$ sending r_1 to r_3, r_3 to r_4, r_4 to r_1, and fixing r_2, ρ be a linear transformation of $\mathcal{V}_4$ into itself arising from σ, then $\rho(\Phi_{D_4}) = \Phi_{D_4}$. The Chevalley group $D_4(q^3)$ has a field automorphism $F_q \colon \mathbb{F}_{q^3} \to \mathbb{F}_{q^3} : t \mapsto t^q$ sending $x_r(t)$ to $x_r(t^q)$, and a graph automorphism ρ sending $x_r(t)$ to $x_{\rho(r)}(t)$ $(r \in \Phi_{D_4})$. Let $F := \rho F_q = F_q \rho$. For a subgroup X of $D_4(q^3)$, set $X^F := \{x \in X | F(x) = x\}$. Then we get $D_4(q^3)^F = {}^3D_4(q^3)$. Set $r^1 := \frac{1}{3}(r + \rho(r) + \rho^2(r))$ for all $r \in \Phi_{D_4}$. Let $\alpha := \frac{1}{3}(r_1 + r_3 + r_4)$ and $\beta := r_2$, then $\Delta_{G_2} := \{\alpha, \beta\}$ (α is short) is a fundamental system of type G_2, and the root system of type G_2 is

$$\Phi_{G_2} = \{r^1 \mid r \in \Phi_{D_4}\} = \{\pm\alpha, \pm\beta, \pm(\alpha+\beta), \pm(2\alpha+\beta), \pm(3\alpha+\beta), \pm(3\alpha+2\beta)\}.$$

We obtain the following elements of ${}^3D_4(q^3)$. Let $r \in \Phi_{D_4}$ and $t \in \mathbb{F}_{q^3}$, we set

$$x_{r^1}(t) := \begin{cases} x_r(t) & \text{if } \rho(r) = r,\ t^q = t \\ x_r(t) x_{\rho(r)}(t^q) x_{\rho^2(r)}(t^{q^2}) & \text{if } \rho(r) \neq r,\ t^{q^3} = t \end{cases}.$$

The root subgroups of ${}^3D_4(q^3)$ are $X_{r^1} := \{x_{r^1}(t) \mid t \in \mathbb{F}_{q^3}\}$ for all $r \in \Phi_{D_4}$.

We write

$$\begin{aligned}
x_1(t) :=& x_{r_1^1}(t) = x_{r_1}(t) x_{r_3}(t^q) x_{r_4}(t^{q^2}) && \text{for all } t \in \mathbb{F}_{q^3}, \\
x_2(t) :=& x_{r_2^1}(t) = x_{r_2}(t) && \text{for all } t \in \mathbb{F}_q, \\
x_3(t) :=& x_{r_5^1}(t) = x_{r_5}(t) x_{r_6}(t^q) x_{r_7}(t^{q^2}) && \text{for all } t \in \mathbb{F}_{q^3}, \\
x_4(t) :=& x_{r_8^1}(t) = x_{r_8}(t) x_{r_{10}}(t^q) x_{r_9}(t^{q^2}) && \text{for all } t \in \mathbb{F}_{q^3}, \\
x_5(t) :=& x_{r_{11}^1}(t) = x_{r_{11}}(t) && \text{for all } t \in \mathbb{F}_q, \\
x_6(t) :=& x_{r_{12}^1}(t) = x_{r_{12}}(t) && \text{for all } t \in \mathbb{F}_q.
\end{aligned}$$

Set $X_i := \{x_i(t) \mid t \in \mathbb{F}_{q^3}\}$ $(i = 1, 3, 4)$ and $X_i := \{x_i(t) \mid t \in \mathbb{F}_q\}$ $(i = 2, 5, 6)$. Note that the root subgroups are denoted by Y_i in [Le13], and $Y_1 := X_2$, $Y_2 := X_1$ and $Y_i := X_i$ for $i = 3, 4, 5, 6$. We obtain a Sylow p-subgroup of ${}^3D_4(q^3)$:

$${}^3D_4^{syl}(q^3) := \left\{ x_2(t_2) x_1(t_1) x_3(t_3) x_4(t_4) x_5(t_5) x_6(t_6) \,\middle|\, t_1, t_3, t_4 \in \mathbb{F}_{q^3},\ t_2, t_5, t_6 \in \mathbb{F}_q \right\},$$

where the product can be taken in an arbitary, but fixed, order.

Let $t_1, t_3, t_4 \in \mathbb{F}_{q^3}$, $t_2, t_5, t_6 \in \mathbb{F}_q$ and define the commutator

$$[x_i(t_i), x_j(t_j)] := x_i(t_i)^{-1} x_j(t_j)^{-1} x_i(t_i) x_j(t_j).$$

Then the non-trivial commutators of ${}^3D_4^{syl}(q^3)$ are determined as follows

$$\begin{aligned}
[x_1(t_1), x_2(t_2)] =& x_3(-t_2 t_1) x_4(t_2 t_1^{q+1}) x_5(-t_2 t_1^{q^2+q+1}) x_6(2 t_2^2 t_1^{q^2+q+1}), \\
[x_1(t_1), x_3(t_3)] =& x_4(t_1 t_3^q + t_1^q t_3) x_5(-t_1^{q+1} t_3^{q^2} - t_1^{q^2+q} t_3 - t_1^{q^2+1} t_3^q) x_6(-t_1 t_3^{q^2+q} - t_1^q t_3^{q^2+1} - t_1^{q^2} t_3^{q+1}), \\
[x_1(t_1), x_4(t_4)] =& x_5(t_1 t_4^q + t_1^q t_4^{q^2} + t_1^{q^2} t_4), \\
[x_3(t_3), x_4(t_4)] =& x_6(t_3 t_4^q + t_3^q t_4^{q^2} + t_3^{q^2} t_4), \\
[x_2(t_2), x_5(t_5)] =& x_6(t_2 t_5).
\end{aligned}$$

From now on, we set

$$x(t_1,t_2,t_3,t_4,t_5,t_6) := x_2(t_2)x_1(t_1)x_3(t_3)x_4(t_4)x_5(t_5)x_6(t_6) \in {}^3D_4^{syl}(q^3).$$

3 Conjugacy classes

In this section, we determine the conjugacy classes of ${}^3D_4^{syl}(q^3)$. Let $U := {}^3D_4^{syl}(q^3)$.

3.1 Notation. *Let $x, u \in U$, then the conjugate of x by u is defined by ${}^u x := uxu^{-1}$, and the conjugacy class of x is the set ${}^U x := \{vxv^{-1} \mid v \in U\}$.*

3.2 Lemma. *Let $\phi_0 \colon \mathbb{F}_{q^3} \to \mathbb{F}_q : t \mapsto t + t^q + t^{q^2}$, then ϕ_0 is an $\mathbb{F}_q$-epimorphism and $|\ker \phi_0| = q^2$.*

Proof. Let $t, s \in \mathbb{F}_{q^3}$ and $k \in \mathbb{F}_q$, then $\phi_0(t+s) = (t+s) + (t+s)^q + (t+s)^{q^2} = (t + t^q + t^{q^2}) + (s + s^q + s^{q^2}) = \phi_0(t) + \phi_0(s)$, and $\phi_0(kt) = (kt) + (kt)^q + (kt)^{q^2} = k(t + t^q + t^{q^2}) = k\phi_0(t)$. Thus ϕ_0 is an $\mathbb{F}_q$-homomorphism. Observe that $\ker \phi_0 = \{t \in \mathbb{F}_{q^3} \mid t + t^q + t^{q^2} = 0\}$. The degree of $t + t^q + t^{q^2}$ is q^2, so $|\ker \phi_0| \leq q^2$. On the other hand, $|\operatorname{im} \phi_0| \leq q$ since $\operatorname{im} \phi_0 \subseteq \mathbb{F}_q$. Then $|\ker \phi_0| = \frac{|\mathbb{F}_{q^3}|}{|\operatorname{im} \phi_0|} \geq q^2$ since $\mathbb{F}_{q^3}/\ker \phi_0 \cong \operatorname{im} \phi_0$. Thus $|\ker \phi_0| = q^2$ and $\operatorname{im} \phi_0 = \mathbb{F}_q$. Therefore, ϕ_0 is an $\mathbb{F}_q$-epimorphism. □

3.3 Proposition. *There exists an element $\eta \in \mathbb{F}_{q^3} \backslash \mathbb{F}_q$ (i.e. $\eta \in \mathbb{F}_{q^3}$ but $\eta \notin \mathbb{F}_q$) such that $\eta^{q^2} + \eta^q + \eta = 1$.*

Proof. By 3.2, we obtain $\#\{t \in \mathbb{F}_{q^3} \mid t^{q^2} + t^q + t = 1\} = q^2$. But $|\mathbb{F}_q| = q$, so there are at least $q^2 - q$ (> 1) elements of $\mathbb{F}_{q^3}$ which satisfy the equation $t^{q^2} + t^q + t = 1$. Then the claim is proved. □

From now on, we fix an element $\eta \in \mathbb{F}_{q^3} \backslash \mathbb{F}_q$ such that $\eta^{q^2} + \eta^q + \eta = 1$.

3.4 Corollary. $1 + \eta^{1-q^2} \neq 0$.

Proof. Suppose that $1 + \eta^{1-q^2} = 0$. Multiply η^{q^2} $(\neq 0)$ to both sides of the formula, then $\eta^{q^2} + \eta = 0$, by 3.3 $\eta^q = 1$, so $\eta = 1 \in \mathbb{F}_q$. This is a contradiction to that $\eta \in \mathbb{F}_{q^3} \backslash \mathbb{F}_q$. □

3.5 Notation/Lemma. *The map $\pi_q \colon \mathbb{F}_{q^3} \to \mathbb{F}_q : x \mapsto \phi_0(\eta x) = (\eta x)^{q^2} + (\eta x)^q + \eta x$ is an $\mathbb{F}_q$-epimorphism and $\mathbb{F}_{q^3} = \ker \pi_q \oplus \mathbb{F}_q$. In particular, $\pi_q|_{\mathbb{F}_q} = \mathrm{id}_{\mathbb{F}_q}$ and $\pi_q^2 = \pi_q$.*

3.6 Lemma. *Let $u \in \mathbb{F}_{q^3}^*$, then the map $\zeta_u \colon \mathbb{F}_{q^3} \to \mathbb{F}_{q^3} : t \mapsto ut^{q^2} + u^q t^q$ is an $\mathbb{F}_q$-automorphism.*

Proof. Since taking the q-th power is an $\mathbb{F}_q$-homomorphism, ζ_u is an $\mathbb{F}_q$-homomorphism. Let $t \in \ker \zeta_u$, then $0 = \zeta_u(t) = ut^{q^2} + u^q t^q = (u^{1-q}t^{q^2-q} + 1)u^q t^q = ((u^{-1}t^q)^{q-1} + 1)u^q t^q$. Suppose that $t \neq 0$, then $(u^{-1}t^q)^{q-1} = -1$ and $(u^{-1}t^q)^{2(q-1)} = 1$. Thus the order of $(u^{-1}t^q)^{q-1}$ is 2, i.e. $|(u^{-1}t^q)^{q-1}| = 2$. On the other hand, since $u^{-1}t^q \in \mathbb{F}_{q^3}^*$, the order $|u^{-1}t^q|$ divides $(q^3 - 1) = (q-1)(q^2+q+1)$, so $|(u^{-1}t^q)^{q-1}|$ divides $(q^2 + q + 1)$. Since $q^2 + q + 1$ is odd, we get $|(u^{-1}t^q)^{q-1}| \neq 2$. This is a contradiction, so $\ker \zeta_u = \{0\}$. Thus $\mathbb{F}_{q^3} \cong \operatorname{im} \zeta_u \subseteq \mathbb{F}_{q^3}$, then $\operatorname{im} \zeta_u = \mathbb{F}_{q^3}$. Therefore, ζ_u is an $\mathbb{F}_q$-automorphism. □

3.7 Corollary. *The map $\mathbb{F}_{q^3} \to \mathbb{F}_{q^3} : t \mapsto t + t^q$ is an $\mathbb{F}_q$-automorphism.*

We obtain the following conjugate elements from the commutator relations of U.

3.8 Lemma. *Let $u := x(r_1, r_2, r_3, r_4, r_5, r_6) \in U$, $x_i(t_i) \in U$ and $\bar{t}_3 \in \mathbb{F}_{q^3}$, then*

$$\begin{aligned}
{}^u x_6(t_6) =& x_6(t_6),\\
{}^u x_5(t_5) =& x_5(t_5)\cdot x_6(r_2t_5),\\
{}^u x_4(t_4) =& x_4(t_4)\cdot x_5(\phi_0\big(r_1t_4^q\big))\cdot x_6(\phi_0\big(r_1r_2t_4^q\big)+\phi_0\big(r_3t_4^q\big)),\\
{}^u x_3(t_3) =& x_3(t_3)\cdot x_4(r_1t_3^q+r_1^qt_3)\cdot x_5(\phi_0\big(r_1^{q^2+q}t_3\big))\\
&\cdot x_6(\phi_0\big(r_1^{q^2+q}r_2t_3\big)+\phi_0\big(-r_1t_3^{q^2+q}\big)+\phi_0\big(-t_3r_4^q\big)),\\
{}^u x_2(t_2) =& x_2(t_2)\cdot x_3(-r_1t_2)\cdot x_4(-t_2r_1^{q+1})\cdot x_5(-t_2r_1^{q^2+q+1})\\
&\cdot x_6(-t_2r_5-t_2^2r_1^{q^2+q+1}-t_2r_1^{q^2+q+1}r_2),\\
{}^u x_1(t_1) =& x_1(t_1)\cdot x_3(r_2t_1)\cdot x_4(-r_2t_1^{q+1}-t_1r_3^q-t_1^qr_3)\\
&\cdot x_5(r_2t_1^{q^2+q+1}+\phi_0\big(r_1^{q^2}(-r_3t_1^q-r_3^qt_1)\big)+\phi_0\big(r_3^{q^2}t_1^{q+1}\big)+\phi_0\big(-t_1r_4^q\big))\\
&\cdot x_6(2r_2^2t_1^{q^2+q+1}+\phi_0\big(r_1^{q^2}r_2(-r_3t_1^q-r_3^qt_1)\big)+\phi_0\big(r_2r_3^{q^2}t_1^{q+1}\big)\\
&\quad+\phi_0\big(-r_2r_4^qt_1\big)+\phi_0\big(-t_1r_3^{q^2+q}\big)),
\end{aligned}$$

and

$$\begin{aligned}
{}^u\big(x_3(t_3)x_5(t_5)\big) =& x_3(t_3)\cdot x_4(r_1t_3^q+r_1^qt_3)\cdot x_5(t_5+\phi_0\big(r_1^{q^2+q}t_3\big))\\
&\cdot x_6(r_2t_5+\phi_0\big(r_1^{q^2+q}r_2t_3\big)+\phi_0\big(-r_1t_3^{q^2+q}\big)+\phi_0\big(-t_3r_4^q\big)),
\end{aligned}$$

$$\begin{aligned}
{}^u\big(x_2(t_2)x_4(t_4)x_5(t_5)\big) =& x_2(t_2)\cdot x_3(-r_1t_2)\cdot x_4(t_4-t_2r_1^{q+1})\cdot x_5(t_5-t_2r_1^{q^2+q+1}+\phi_0\big(r_1t_4^q\big))\\
&\cdot x_6(-t_2r_5-t_2^2r_1^{q^2+q+1}-t_2r_1^{q^2+q+1}r_2+\phi_0\big(r_1r_2t_4^q\big)+\phi_0\big(r_3t_4^q\big)+r_2t_5),
\end{aligned}$$

$$\begin{aligned}
&{}^u\big(x_1(t_1)x_3(\bar{t}_3)\big)\\
=&x_1(t_1)\cdot x_3(r_2t_1+\bar{t}_3)\cdot x_4(-r_2t_1^{q+1}-t_1r_3^q-t_1^qr_3+r_1\bar{t}_3^q+r_1^q\bar{t}_3)\\
&\cdot x_5(r_2t_1^{q^2+q+1}+\phi_0\big(r_1^{q^2}(-r_3t_1^q-r_3^qt_1)\big)+\phi_0\big(r_3^{q^2}t_1^{q+1}\big)+\phi_0\big(-t_1r_4^q\big)+\phi_0\big(r_1^{q^2+q}\bar{t}_3\big))\\
&\cdot x_6(2r_2^2t_1^{q^2+q+1}+\phi_0\big(r_1^{q^2}r_2(-r_3t_1^q-r_3^qt_1)\big)+\phi_0\big(r_2r_3^{q^2}t_1^{q+1}\big)+\phi_0\big(-r_2r_4^qt_1\big)+\phi_0\big(-t_1r_3^{q^2+q}\big)\\
&\quad+\phi_0\big(r_1^{q^2+q}r_2\bar{t}_3\big)+\phi_0\big(-r_1\bar{t}_3^{q^2+q}\big)+\phi_0\big(-\bar{t}_3r_4^q\big)+\phi_0\big(\bar{t}_3^{q^2}(r_2t_1^{q+1}+t_1r_3^q+t_1^qr_3)\big)),
\end{aligned}$$

$$\begin{aligned}
&{}^u\big(x_2(t_2)x_1(t_1)\big)\\
=&x_2(t_2)x_1(t_1)\cdot x_3(r_2t_1-r_1t_2)\cdot x_4(-r_2t_1^{q+1}-t_1r_3^q-t_1^qr_3-t_2r_1^{q+1}+t_1t_2r_1^q+t_1^qt_2r_1)\\
&\cdot x_5(r_2t_1^{q^2+q+1}+\phi_0\big(r_1^{q^2}(-r_3t_1^q-r_3^qt_1)\big)+\phi_0\big(r_3^{q^2}t_1^{q+1}\big)+\phi_0\big(-t_1r_4^q\big)\\
&\quad-t_2r_1^{q^2+q+1}+\phi_0\big(-r_1t_1^{q^2+q}t_2\big)+\phi_0\big(t_1^{q^2}t_2r_1^{q+1}\big))\\
&\cdot x_6(2r_2^2t_1^{q^2+q+1}+\phi_0\big(r_1^{q^2}r_2(-r_3t_1^q-r_3^qt_1)\big)+\phi_0\big(r_2r_3^{q^2}t_1^{q+1}\big)+\phi_0\big(-r_2r_4^qt_1\big)+\phi_0\big(-t_1r_3^{q^2+q}\big)\\
&\quad-t_2r_5-t_2^2r_1^{q^2+q+1}-t_2r_1^{q^2+q+1}r_2+\phi_0\big(-2r_1r_2t_1^{q^2+q}t_2\big)+\phi_0\big(t_1^{q^2}t_2r_1^{q+1}r_2\big)\\
&\quad+\phi_0\big(r_1^{q^2+q}t_1t_2^2\big)).
\end{aligned}$$

Before we get the conjugacy classes of ${}^3D_4^{syl}(q^3)$, we need the following lemma.

3.9 Lemma. *Let $u \in \mathbb{F}_{q^3}\backslash\mathbb{F}_q$, $\ker\phi_0$ be the kernel of ϕ_0, $S := u\ker\phi_0$ and*

$$\phi_0|_S\colon u\ker\phi_0 \to \mathbb{F}_q : t \mapsto t^{q^2}+t^q+t.$$

Then $\phi_0|_S$ is an $\mathbb{F}_q$-epimorphism.

Proof. The kernel $\ker\phi_0$ is an $\mathbb{F}_q$-vector subspace of $\mathbb{F}_{q^3}$ and $u \in \mathbb{F}_{q^3}$, so $u\ker\phi_0$ is an $\mathbb{F}_q$-vector subspace of $\mathbb{F}_{q^3}$. Thus $\phi_0|_S$ is an $\mathbb{F}_q$-homomorphism by 3.2. Let $t_0 \in \ker\phi_0$ and $ut_0 \in \ker(\phi_0|_S)$, then $t_0^{q^2}+t_0^q+t_0=0$ and $u^{q^2}t_0^{q^2}+u^qt_0^q+ut_0=0$, so $u^{q^2}t_0^{q^2}+u^{q^2}t_0^q+u^{q^2}t_0=0$ since $u^{q^2}\neq 0$. Then $(u^q-u^{q^2})t_0^q+(u-u^{q^2})t_0=0$. Since $u\notin\mathbb{F}_q$, $u^q-u^{q^2}\neq 0$. Then $(u^q-u^{q^2})^{-1}t_0=(u^q-u^{q^2})^{-q}t_0^q$, so $(u^q-u^{q^2})^{-1}t_0\in\mathbb{F}_q$ and $t_0\in(u^q-u^{q^2})\mathbb{F}_q$. Thus $|\ker(\phi_0|_S)|\le|x\mathbb{F}_q|=q$, then $|\mathrm{im}(\phi_0|_S)|\ge\frac{|S|}{|\ker(\phi_0|_S)|}\overset{3.2}{=}\frac{q^2}{q}=q$. We have $|\mathrm{im}(\phi_0|_S)|=q$ since $|\mathrm{im}(\phi_0|_S)|\le|\mathbb{F}_q|=q$. Therefore, $\phi_0|_S$ is an $\mathbb{F}_q$-epimorphism. □

3.10 Notation. *Let $a^*\in\mathbb{F}_{q^3}^*=\mathbb{F}_{q^3}\backslash\{0\}$, then denote by T^{a^*} a complete set of coset representatives (i.e. a transversal) of $(a^*\mathbb{F}_q^+)$ in $\mathbb{F}_{q^3}^+$. Thus $|T^{a^*}|=q^2$. If $\bar{t}_0\in T^{a^*}$ and $\bar{t}_0\in a^*\mathbb{F}_q^+$, we set $\bar{t}_0=0$.*

3.11 Proposition (Conjugacy classes of ${}^3D_4^{syl}(q^3)$)**.** *The conjugacy classes of ${}^3D_4^{syl}(q^3)$ are listed in Table 1.*

Table 1: Conjugacy classes of ${}^3D_4^{syl}(q^3)$

Representatives x	$\#\{{}^Ux\}$	${}^Ux$	$\lvert{}^Ux\rvert$
I_8	1	$x(0,0,0,0,0,0)$	1
$x_6(t_6^*)$	$q-1$	$x(0,0,0,0,0,t_6^*)$	1
$x_5(t_5^*)$	$q-1$	$x(0,0,0,0,t_5^*,s_6)$ $s_6\in\mathbb{F}_q$	q
$x_4(t_4^*)$	q^3-1	$x(0,0,0,t_4^*,s_5,s_6)$ $s_5,s_6\in\mathbb{F}_q$	q^2
$x(0,0,t_3^*,0,t_5,0)$	$(q^3-1)q$	$x(0,0,t_3^*,s_4,\hat{s}_5,s_6)$ $s_4\in\mathbb{F}_{q^3},s_6\in\mathbb{F}_q$	q^4
$x(0,t_2^*,0,t_4,t_5,0)$	$(q-1)q^4$	$x(0,t_2^*,s_3,\hat{s}_4,\hat{s}_5,s_6)$ $s_3\in\mathbb{F}_{q^3},s_6\in\mathbb{F}_q$	q^4
$x(t_1^*,0,0,0,0,t_6)$	$(q^3-1)q$	$x(t_1^*,0,t_1^*s_2,s_4,s_5,\hat{s}_6)$ $s_4\in\mathbb{F}_{q^3},s_2,s_5\in\mathbb{F}_q$	q^5
$x(t_1^*,0,\bar{t}_3^*,0,0,0)$	$(q^3-1)(q^2-1)$	$x(t_1^*,0,\bar{t}_3^*+t_1^*s_2,s_4,s_5,s_6)$ $s_4\in\mathbb{F}_{q^3},s_2,s_5,s_6\in\mathbb{F}_q$	q^6
$x(t_1^*,t_2^*,0,0,0,0)$	$(q-1)(q^3-1)$	$x(t_1^*,t_2^*,s_3,s_4,s_5,s_6)$ $s_3,s_4\in\mathbb{F}_{q^3},s_5,s_6\in\mathbb{F}_q$	q^8

where $0\neq\bar{t}_3^*\in T^{t_1^*}$, $\hat{s}_-$ is determined by some of t_-^*, t_- and s_-.

Proof. We prove the hard cases of the proposition. Let $0\neq t_1\in\mathbb{F}_{q^3}^*$, $\bar{t}_3\in\mathbb{F}_{q^3}$, $t_6\in\mathbb{F}_q$, $u:=x(r_1,r_2,r_3,r_4,r_5,r_6)\in U$ and $x(a_1,a_2,a_3,a_4,a_5,a_6):={}^u\big(x_1(t_1)x_3(\bar{t}_3)x_6(t_6)\big)$.

(1) Let $\bar{t}_3 \in t_1\mathbb{F}_q^+$, then there exists $s \in \mathbb{F}_q^+$ such that $\bar{t}_3 = st_1$. By 3.8,

$$\begin{aligned}
a_1 =& t_1,\ a_2 = 0,\ a_3 = r_2t_1 + \bar{t}_3 = (r_2+s)t_1,\ a_4 = -r_2t_1^{q+1} - (r_3 - sr_1)t_1^q - (r_3 - sr_1)^q t_1,\\
a_5 =& r_2t_1^{q^2+q+1} + \phi_0\big(r_1^{q^2}(-r_3t_1^q - r_3^q t_1)\big) + \phi_0\big(r_3^{q^2}t_1^{q+1}\big) + \phi_0\big(-t_1r_4^q\big) + \phi_0\big(sr_1^{q^2+q}t_1\big),\\
a_6 =& t_6 + r_2^2t_1^{q^2+q+1} + r_2a_5 + \phi_0\big(-t_1r_3^{q^2+q}\big) + \phi_0\big(-s^2r_1t_1^{q^2+q}\big)\\
&+ \phi_0\big(-st_1r_4^q\big) + \phi_0\big(st_1^{q^2}(r_2t_1^{q+1} + t_1r_3^q + t_1^q r_3)\big)\\
=& t_6 + r_2^2t_1^{q^2+q+1} + (r_2+s)a_5 + 2sr_2t_1^{q^2+q+1}\\
&+ \phi_0\big(s(r_3 - sr_1)t_1^{q^2+q}\big) - \phi_0\big((r_3 - sr_1)^{q^2+q}t_1\big).
\end{aligned}$$

Let a_3, a_4 and a_5 be fixed, then a_6 are determined uniquely 3.2 and 3.6. Hence we get the conjugacy classes of $x_1(t_1)x_6(t_6)$.

$$^U\big(x_1(t_1)x_6(t_6)\big) = \big\{x(t_1, 0, s_2t_1, s_4, s_5, \hat{s}_6) \mid s_4 \in \mathbb{F}_{q^3}, s_2, s_5 \in \mathbb{F}_q\big\}.$$

(2) Let $\bar{t}_3 \notin t_1\mathbb{F}_q^+$. By 3.8,

$$\begin{aligned}
a_1 =& t_1, \quad a_2 = 0, \quad a_3 = \bar{t}_3 + r_2t_1, \quad a_4 = -r_2t_1^{q+1} - t_1r_3^q - t_1^q r_3 + r_1\bar{t}_3^q + r_1^q\bar{t}_3,\\
a_5 =& r_2t_1^{q^2+q+1} + \phi_0\big(r_1^{q^2}(-r_3t_1^q - r_3^q t_1)\big) + \phi_0\big(r_3^{q^2}t_1^{q+1}\big) + \phi_0\big(-t_1r_4^q\big) + \phi_0\big(r_1^{q^2+q}\bar{t}_3\big),\\
a_6 =& t_6 + r_2^2t_1^{q^2+q+1} + r_2a_5 + \phi_0\big(-t_1r_3^{q^2+q}\big) + \phi_0\big(-r_1\bar{t}_3^{q^2+q}\big)\\
&+ \phi_0\big(-\bar{t}_3r_4^q\big) + \phi_0\big(\bar{t}_3^{q^2}(r_2t_1^{q+1} + t_1r_3^q + t_1^q r_3)\big).
\end{aligned}$$

For the fixed r_1, r_2, r_3 and a_5, let

$$T := \{t_1r_4^q \mid r_4 \in \mathbb{F}_{q^3}, \text{and } r_1, r_2, r_3, a_5 \text{ are fixed}\}.$$

By 3.2, $|T| = q^2$. Let $x_0 \in T$, then $T = x_0 + \ker\phi_0$ and $\bar{t}_3r_4^q \in \frac{\bar{t}_3x_0}{t_1} + \frac{\bar{t}_3}{t_1}\ker\phi_0$. We know $\bar{t}_3 \notin t_1\mathbb{F}_q^+$, so $\frac{\bar{t}_3}{t_1} \in \mathbb{F}_{q^3}^+ \backslash \mathbb{F}_q^+$. Hence

$$\{\phi_0(\bar{t}_3r_4^q) \mid t_1r_4^q \in T\} = \left\{\phi_0(\frac{\bar{t}_3x_0}{t_1}) + \phi_0(t) \,\middle|\, t \in \frac{\bar{t}_3}{t_1}\ker\phi_0\right\} \overset{3.9}{=} \mathbb{F}_q^+.$$

Thus a_6 can be every element of $\mathbb{F}_q$ for the fixed r_1, r_2, r_3 and a_5. Therefore,

$$^U\big(x_1(t_1)x_3(\bar{t}_3)\big) = \big\{x(t_1, 0, \bar{t}_3 + s_2t_1, s_4, s_5, s_6) \mid s_4 \in \mathbb{F}_{q^3}, s_2, s_5, s_6 \in \mathbb{F}_q\big\}.$$

By 3.8, the other conjugacy classes are also determined. Then the proposition is obtained. □

4 Irreducible characters

In [Le13], Tung Le constructed and counted all ordinary irreducible characters of ${}^3D_4^{syl}(q^3)$, mainly using Clifford theory. In this section, we give the specific constructions of the irreducible characters of ${}^3D_4^{syl}(q^3)$ using Clifford's Theorem (see [CR81]), so that we can determine the values of the irreducible characters in the next section.

Let G be a finite group, N a normal subgroup of G, K a field, and $U := {}^3D_4^{syl}(q^3)$. Let $\mathrm{Irr}(G)$ be the set of all complex irreducible characters of G, and triv_G the trivial character of G.

Let H be a subgroup of G, $\chi \in \mathrm{Irr}(G)$ and $\lambda \in \mathrm{Irr}(H)$, then we denote by $\mathrm{Ind}_H^G\lambda$ the character induced from λ, and deonte by $\mathrm{Res}_H^G\chi$ the restriction of χ to H. The center of G is denoted by $Z(G)$. The kernel of χ is $\ker\chi := \{g \in G \mid \chi(g) = \chi(1)\}$. The commutator subgroup of G is $G' = \langle\, [x,y] \mid x, y \in G \,\rangle$, where $[x,y] = x^{-1}y^{-1}xy$. Let $\lambda \in \mathrm{Irr}(N)$, then the inertia group in G is $I_G(\lambda) = \{g \in G \mid \lambda^g = \lambda\}$ where $\lambda^g(n) = \lambda(gng^{-1})$ for all $n \in N$. In paricular, $N \trianglelefteq I_G(\lambda) \leqslant G$.

4.1 Notation/Lemma. *Let $\vartheta\colon \mathbb{F}_q^+ \to \mathbb{C}^*$ denote a fixed nontrivial linear character of the additive group $\mathbb{F}_q^+$ of $\mathbb{F}_q$ once and for all. In particular, $\sum_{x\in\mathbb{F}_q^+}\vartheta(x) = 0$.*

We determine the irreducible characters of $\mathbb{F}_{q^3}^+$ and $\mathbb{F}_q^+$.

4.2 Lemma. *Let $a \in \mathbb{F}_{q^3}$, $b \in \mathbb{F}_q$ and*

$$\vartheta_a\colon \mathbb{F}_{q^3}^+ \to \mathbb{C}^* : x \mapsto \vartheta\pi_q(ax),$$
$$\vartheta_b\colon \mathbb{F}_q^+ \to \mathbb{C}^* : y \mapsto \vartheta\pi_q(by) = \vartheta(by),$$

then $\mathrm{Irr}(\mathbb{F}_{q^3}^+) = \{\vartheta_a \mid a \in \mathbb{F}_{q^3}\}$ and $\mathrm{Irr}(\mathbb{F}_q^+) = \{\vartheta_b \mid b \in \mathbb{F}_q\}$.

Proof. Let $a \in \mathbb{F}_{q^3}$. By 3.5 and 4.1, we get $\vartheta_a(x+y) = \vartheta_a(x)\cdot\vartheta_a(y)$ for all $x, y \in \mathbb{F}_{q^3}$, so $\mathrm{Irr}(\mathbb{F}_{q^3}^+) \supseteq \{\vartheta_a \mid a \in \mathbb{F}_{q^3}\}$. Let $a, c \in \mathbb{F}_{q^3}$ and $\vartheta_a = \vartheta_c$, then $\vartheta_a(x) = \vartheta_c(x)$ for all $x \in \mathbb{F}_{q^3}$, i.e. $\vartheta\pi_q(ax) = \vartheta\pi_q(cx)$ for all $x \in \mathbb{F}_{q^3}$. Thus $a = c$ by 3.5. Otherwise, $\vartheta(y) = 1$ for all $y \in \mathbb{F}_{q^3}$, this is a contradiction since ϑ is nontrivial by 4.1. Thus $\#\{\vartheta_a \mid a \in \mathbb{F}_{q^3}\} = q^3$. Therefore, $\mathrm{Irr}(\mathbb{F}_{q^3}^+) = \{\vartheta_a \mid a \in \mathbb{F}_{q^3}\}$. Similarly, we obtain the second formula. □

4.3 Corollary. *Let $u \in \mathbb{F}_{q^3}$, and $\tilde{\vartheta}_u\colon \mathbb{F}_{q^3}^+ \to \mathbb{C}^* : t \mapsto \vartheta\pi_q(u^qt + ut^{q^2})$, then $\tilde{\vartheta}_u = \vartheta_{\eta^{-1}(\eta+\eta^q)u^q}$ and $\mathrm{Irr}(\mathbb{F}_{q^3}^+) = \{\tilde{\vartheta}_u \mid u \in \mathbb{F}_{q^3}\}$.*

Proof. For all $t \in \mathbb{F}_{q^3}^+$, $\tilde{\vartheta}_u(t) = \vartheta\pi_q(u^qt + ut^{q^2}) \overset{3.5}{=} \vartheta\phi_0(\eta u^qt + \eta ut^{q^2}) \overset{3.2}{=} \vartheta\phi_0((\eta+\eta^q)u^qt)$ $=\vartheta\pi_q(\eta^{-1}(\eta+\eta^q)u^qt) = \vartheta_{\eta^{-1}(\eta+\eta^q)u^q}(t)$. By 3.4, $\eta + \eta^q \neq 0$. By 4.2, we obtain the second formula. □

4.4 Lemma. *Let G be a finite group, $Z(G) \subseteq N \trianglelefteq G$, and $\chi \in \mathrm{Irr}(G)$. Let $\lambda \in \mathrm{Irr}(N)$ such that $\langle \mathrm{Res}_N^G\chi, \lambda\rangle_N = e > 0$. Then for all $g \in Z(G)$,*

$$\left(\mathrm{Res}_N^G\chi\right)(g) = e\frac{|G|}{|I_G(\lambda)|}\lambda(g)$$

and $g \notin \ker\chi \iff g \notin \ker\lambda$. In particular, if $X \leqslant Z(G)$, then $X \nsubseteq \ker\chi$ if and only if $X \nsubseteq \ker\lambda$.

Proof. By Clifford's Theorem, we have for all $g \in Z(G)$

$$\left(\mathrm{Res}_N^G\right)\chi(g) = e\sum_{h\in T_{I_G(\lambda)\backslash G}}\lambda^h(g) \overset{g\in Z(G)}{=} e\frac{|G|}{|I_G(\lambda)|}\lambda(g).$$

Let 1 be the identity element of G and $g_0 \in Z(G)$, then

$$\chi(g_0) \neq \chi(1) \iff \left(\mathrm{Res}_N^G\chi\right)(g_0) \neq \left(\mathrm{Res}_N^G\chi\right)(1) \iff \lambda(g_0) \neq \lambda(1).$$

Thus $g \notin \ker\chi \iff g \notin \ker\lambda$. □

4.5 Definition (2.1, [Le13]). *Let χ be an irreducible character of a group G. χ is said to be **almost faithful** if $Z(G) \not\leqslant \ker\chi$.*

We determine some inertia groups.

4.6 Lemma. *Let $T := X_2X_3X_4X_5X_6$, $N := X_4X_5X_6$ and $H := X_1X_4X_5X_6$.*

(1) The subgroup N is abelian, $N \trianglelefteq U$, $T \trianglelefteq U$ and $H \leqslant U$ as follows



(2) Let $\lambda \in \mathrm{Irr}(N)$ and $\mathrm{Res}^N_{X_6}\lambda \neq \mathrm{triv}_{X_6}$. If λ satisfies that $\mathrm{Res}^N_{X_5}\lambda = \mathrm{triv}_{X_5}$, we have

$$I_U(\lambda) = \{u \in U \mid \lambda^u = \lambda\} = H.$$

(3) Let $\lambda \in \mathrm{Irr}(N)$, the inertia group $I_T(\lambda)$ is

$$I_T(\lambda) = \begin{cases} T & \text{if } \mathrm{Res}^N_{X_6}\lambda = \mathrm{triv}_{X_6} \\ N & \text{if } \mathrm{Res}^N_{X_6}\lambda \neq \mathrm{triv}_{X_6} \end{cases}.$$

(4) Let $\lambda \in \mathrm{Irr}(N)$, then the inertia group $I_H(\lambda)$ is

$$I_H(\lambda) = \begin{cases} H & \text{if } \mathrm{Res}^N_{X_5}\lambda = \mathrm{triv}_{X_5} \\ N & \text{if } \mathrm{Res}^N_{X_5}\lambda \neq \mathrm{triv}_{X_5} \end{cases}.$$

(5) Let $\psi \in \mathrm{Irr}(T)$ and $X_6 = Z(T) \nsubseteq \ker\psi$, then the inertia group $I_U(\psi)$ is

$$I_U(\psi) = \{u \in U \mid \psi^u = \psi\} = U.$$

Proof. (1) From the commutator relations of U and by 3.8, we obtain that N is abelian, N and T are normal subgroups of U, and H is a subgroup of U. Then N is a normal subgroup of N and H, since N is a normal subgroup of U.

(2) Let $u := x(r_1, r_2, r_3, r_4, r_5, r_6) \in U$, $x := x_4(t_4)x_5(t_5)x_6(t_6) \in N$, and $\lambda \in \mathrm{Irr}(N)$, then λ is a linear character since N is abelian. We have

$$\begin{aligned} \lambda^u(x) =& \lambda(u \cdot x \cdot u^{-1}) \\ \overset{3.8}{=}& \lambda\left(x_4(t_4)x_5(t_5 + \phi_0(r_1t_4^q))x_6(t_6 + r_2t_5 + \phi_0(r_1r_2t_4^q) + \phi_0(r_3t_4^q))\right) \\ \overset{\mathrm{Res}^N_{X_5}\lambda = \mathrm{triv}_{X_5}}{=}& \lambda(x_4(t_4)) \cdot \lambda(x_6(t_6 + r_2t_5 + \phi_0(r_1r_2t_4^q) + \phi_0(r_3t_4^q))). \end{aligned}$$

Since $X_6 \nsubseteq \ker\lambda$ and $X_5 \subseteq \ker\lambda$, we get

$$\begin{aligned} I_U(\lambda) &= \{u \in U \mid \lambda^u = \lambda\} = \{u \in U \mid \lambda(u \cdot x \cdot u^{-1}) = \lambda(x) \text{ for all } x \in N\} \\ &\overset{3.2}{=} \{u := x(r_1, r_2, r_3, r_4, r_5, r_6) \in U \mid r_2 = r_3 = 0\} = X_1X_4X_5X_6 = H. \end{aligned}$$

(3) c.f. the proof of (2).

(4) c.f. the proof of (2).

(5) Let $x_6(t_6) \in X_6$. Since $X_6 = Z(T) \trianglelefteq N \trianglelefteq T$, there exists $\lambda \in \mathrm{Irr}(N)$ such that $0 < e = \langle \mathrm{Res}_N^T \psi, \lambda \rangle_N$, then $\left(\mathrm{Res}_N^T \psi\right)(x_6(t_6)) = e\frac{|T|}{|I_T(\lambda)|}\lambda(x_6(t_6))$. By 4.4, $X_6 = Z(T) \nsubseteq \ker \lambda$ since $X_6 = Z(T) \nsubseteq \ker \psi$. Then $\mathrm{Res}_{X_6}^N \lambda \neq \mathrm{triv}_{X_6}$. Then $I_T(\lambda) = N$ by (3), so $\mathrm{Ind}_N^T \lambda \in \mathrm{Irr}(T)$ by Clifford's Theorem. Thus $e = 1$ and $\psi = \mathrm{Ind}_N^T \lambda$.

Let $u := x(s_1, s_2, s_3, s_4, s_5, s_6) \in U$ and $h := x(0, t_2, t_3, t_4, t_5, t_6) \in T$, then

$$\psi(h) = \left(\mathrm{Ind}_N^T \lambda\right)(h) = \frac{1}{|N|} \sum_{\substack{y \in T \\ y \cdot h \cdot y^{-1} \in N}} \lambda(y \cdot h \cdot y^{-1}).$$

– Let $t_2 \neq 0$ or $t_3 \neq 0$, then $\psi(h) = \left(\mathrm{Ind}_N^T \lambda\right)(h) = 0$ and

$$(\psi^u)(h) = \left(\mathrm{Ind}_N^T \lambda\right)^u(h) = \left(\mathrm{Ind}_N^T \lambda\right)(u \cdot h \cdot u^{-1}) \overset{3.8}{=} 0.$$

– Let $t_2 = t_3 = 0$, then by 3.8

$$\begin{aligned}\psi(h) = \left(\mathrm{Ind}_N^T \lambda\right)(h) =& \frac{1}{|N|} \sum_{\substack{y \in T \\ y \cdot h \cdot y^{-1} \in N}} \lambda(y \cdot h \cdot y^{-1}) = \frac{1}{|N|} \sum_{y \in T} \lambda(y \cdot h \cdot y^{-1}) \\ \overset{y := x(0, r_2, r_3, r_4, r_5, r_6)}{=}& \frac{1}{|N|} \sum_{\substack{r_2, r_5, r_6 \in \mathbb{F}_q \\ r_3, r_4 \in \mathbb{F}_{q^3}}} (x_4(t_4) x_5(t_5) x_6(t_6 + r_2 t_5 + \phi_0(r_3 t_4^q))),\end{aligned}$$

and

$$\begin{aligned}&(\psi^u)(h) = \left(\mathrm{Ind}_N^T \lambda\right)^u(h) = \left(\mathrm{Ind}_N^T \lambda\right)(u \cdot h \cdot u^{-1}) \\ =& \left(\mathrm{Ind}_N^T \lambda\right)(x_4(t_4) x_5(t_5 + \phi_0(s_1 t_4^q)) x_6(t_6 + s_2 t_5 + \phi_0(s_1 s_2 t_4^q) + \phi_0(s_3 t_4^q))) \\ =& \frac{1}{|N|} \sum_{\substack{r_2, r_5, r_6 \in \mathbb{F}_q \\ r_3, r_4 \in \mathbb{F}_{q^3}}} \lambda(x_4(t_4) x_5(t_5 + \phi_0(s_1 t_4^q)) x_6(t_6 + s_2 t_5 + \phi_0(s_1 s_2 t_4^q) + \phi_0(s_3 t_4^q)) \\ &\cdot x_6(r_2 t_5 + r_2 \phi_0(s_1 t_4^q) + \phi_0(r_3 t_4^q))).\end{aligned}$$

Since $X_6 \nsubseteq \ker \lambda$, we have for all $u \in U$

$$(\psi^u)(h) = \psi(h) = \begin{cases} q^4 \lambda(x_6(t_6)) & \text{if } t_2 = t_3 = t_4 = t_5 = 0 \\ 0 & \text{otherwise} \end{cases}.$$

Thus $I_U(\psi) = U$. □

We determine irreducible characters of the abelian groups $X_4 X_5 X_6$. From now on, we set $N := X_4 X_5 X_6$.

4.7 Lemma. *Let $A_{17}, A_{16} \in \mathbb{F}_q$, $A_{15} \in \mathbb{F}_{q^3}$ and*

$$\lambda^{A_{17},A_{16},A_{15}}(x_4(t_4)x_5(t_5)x_6(t_6)) := \vartheta(A_{17}t_6)\cdot\vartheta(A_{16}t_5)\cdot\vartheta\pi_q(A_{15}^q t_4 + A_{15}t_4^{q^2}),$$

then $\mathrm{Irr}(N) = \left\{\lambda^{A_{17},A_{16},A_{15}} \,\middle|\, A_{17}, A_{16} \in \mathbb{F}_q, A_{15} \in \mathbb{F}_{q^3}\right\}$.

Now we determine the irreducible characters of the subgroup $H = X_1X_4X_5X_6$ of U.

4.8 Lemma. *Let $H := X_1X_4X_5X_6$ and $\tilde{\chi} \in \mathrm{Irr}(H)$, then $H' = X_5$.*

(1) If $X_5 \subseteq \ker\tilde{\chi}$, let $\bar{H}_{146} := X_5\backslash H \cong \bar{X}_1\bar{X}_4\bar{X}_6$, $\bar{\chi}^{A_{17},A_{15},A_{12}} \in \mathrm{Irr}(\bar{H}_{146})$,

$$\bar{\chi}^{A_{17},A_{15},A_{12}}(\bar{x}_1(t_1)\bar{x}_4(t_4)\bar{x}_6(t_6)) := \vartheta(A_{17}t_6)\cdot\vartheta\pi_q(A_{15}^q t_4 + A_{15}t_4^{q^2})\cdot\vartheta\pi_q(A_{12}t_1).$$

Denote by $\tilde{\chi}^{A_{17},A_{15},A_{12}}$ the lift of $\bar{\chi}^{A_{17},A_{15},A_{12}}$ to H, then

$$\mathrm{Irr}(H)_1 := \{\tilde{\chi} \in \mathrm{Irr}(H) \mid X_5 \subseteq \ker\tilde{\chi}\} = \left\{\tilde{\chi}^{A_{17},A_{15},A_{12}} \in \mathrm{Irr}(H) \,\middle|\, A_{17} \in \mathbb{F}_q, A_{15}, A_{12} \in \mathbb{F}_{q^3}\right\}.$$

(2) If $X_5 \nsubseteq \ker\tilde{\chi}$, we have

$$\mathrm{Irr}(H)_2 := \{\tilde{\chi} \in \mathrm{Irr}(H) \mid X_5 \nsubseteq \ker\tilde{\chi}\} = \left\{\mathrm{Ind}_N^H\lambda^{A_{17},A_{16}^*,0} \,\middle|\, A_{17} \in \mathbb{F}_q, A_{16}^* \in \mathbb{F}_q^*\right\}.$$

Thus, $\mathrm{Irr}(H) = \mathrm{Irr}(H)_1\dot{\cup}\mathrm{Irr}(H)_2$, i.e. H has q^7 linear characters and $(q-1)q$ irreducible characters of degree q^3.

Proof. Let $\tilde{\chi} \in \mathrm{Irr}(H)$. From the commutator relations, $H' = X_5$ and $Z(H) = X_5X_6$.

(1) If $X_5 \subseteq \ker\tilde{\chi}$, $\tilde{\chi}$ is linear. Since $H' = X_5$, all linear characters of H are precisely the lifts to H of the irreducible characters of the abelian quotient group $X_5\backslash H$.

(2) If $X_5 \nsubseteq \ker\tilde{\chi}$, let $\lambda \in \mathrm{Irr}(N)$ and $\langle \mathrm{Res}_N^H\tilde{\chi}, \lambda\rangle_N > 0$. Since $X_5 \trianglelefteq Z(H) \trianglelefteq N \trianglelefteq H$, by 4.4 $X_5 \nsubseteq \ker\lambda$ i.e. $\mathrm{Res}_{X_5}^N\lambda \neq \mathrm{triv}_{X_5}$, then $I_H(\lambda) = N$ by (4) of 4.6. By Clifford's Theorem, $\tilde{\chi} = \mathrm{Ind}_N^H\lambda$ and $\deg(\tilde{\chi}) = q^3$. Then there exists $\lambda^{A_{17},A_{16}^*,A_{15}} \in \mathrm{Irr}(N)$ such that $\lambda = \lambda^{A_{17},A_{16}^*,A_{15}}$. We note that X_1 is a transversal of N in H. Let $r_1 \in \mathbb{F}_{q^3}$, for all $x_4(t_4)x_5(t_5)x_6(t_6) \in N$,

$$\begin{aligned}
&\left(\lambda^{A_{17},A_{16}^*,A_{15}}\right)^{x_1(r_1)}(x_4(t_4)x_5(t_5)x_6(t_6))\\
=&\lambda^{A_{17},A_{16}^*,A_{15}}(x_1(r_1)\cdot x_4(t_4)x_5(t_5)x_6(t_6)\cdot x_1(r_1)^{-1})\\
=&\lambda^{A_{17},A_{16}^*,A_{15}}(x_4(t_4)x_5(t_5+\phi_0(r_1t_4^q))x_6(t_6))\\
=&\vartheta(A_{17}t_6)\cdot\vartheta(A_{16}^*(t_5+\phi_0(r_1t_4^q)))\cdot\vartheta\pi_q(A_{15}^qt_4 + A_{15}t_4^{q^2})\\
=&\vartheta(A_{17}t_6)\cdot\vartheta(A_{16}^*t_5)\cdot\vartheta\pi_q(\eta^{-1}((\eta+\eta^q)A_{15}^q + A_{16}^*r_1^{q^2})t_4).
\end{aligned}$$

By Clifford's Theorem, $\tilde{\chi} = \mathrm{Ind}_N^H\lambda^{A_{17},A_{16}^*,0}$ and

$$\mathrm{Res}_N^H\mathrm{Ind}_N^H\lambda^{A_{17},A_{16}^*,0} = \sum_{r_1\in\mathbb{F}_{q^3}}\left(\lambda^{A_{17},A_{16}^*,A_{15}}\right)^{x_1(r_1)} = \sum_{B_{15}\in\mathbb{F}_{q^3}}\lambda^{A_{17},A_{16}^*,B_{15}}.$$

Thus H has q^7 linear characters and $(q-1)q$ irreducible characters of degree q^3. □

Table 2: Character table of H

$\lvert^H x\rvert$	1	q	q
x	$x_5(t_5)x_6(t_6)$	$x_4(t_4^*)x_6(t_6)$	$x_1(t_1^*)x_4(t_4)x_6(t_6)$
$\tilde{\chi}^{A_{17},A_{15},A_{12}}$	$\vartheta(A_{17}t_6)$	$\vartheta(A_{17}t_6)$ $\cdot\vartheta\pi_q(A_{15}^q t_4^* + A_{15}t_4^{*q^2})$	$\vartheta(A_{17}t_6)$ $\cdot\vartheta\pi_q(A_{15}^q t_4 + A_{15}t_4^{q^2})$ $\cdot\vartheta\pi_q(A_{12}t_1^*)$
$\mathrm{Ind}_N^H\lambda^{A_{17},A_{16}^*,0}$	$q^3\vartheta(A_{17}t_6)$ $\cdot\vartheta(A_{16}^*t_5)$	0	0

By the commutator relations, we determine the conjugacy classes of H, and obtain the character table of H.

4.9 Corollary. *Let $x := x(t_1,0,0,t_4,t_5,t_6) \in H$ be a representative of one conjugacy class of H, then the character table of H is shown in Table 2.*

Similar to H, we construct the irreducible character of the normal subgroup $X_2X_3X_4X_5X_6$ of U.

4.10 Lemma. *Let $T := X_2X_3X_4X_5X_6$ and $\psi \in \mathrm{Irr}(T)$, then $T' = X_6$.*

(1) If $X_6 \subseteq \ker\psi$, let $\bar{H}_{2345} := X_6\backslash T \cong \bar{X}_2\bar{X}_3\bar{X}_4\bar{X}_5$, $\bar{\chi}^{A_{16},A_{15},A_{13},A_{23}} \in \mathrm{Irr}(\bar{H}_{2345})$,

$$\begin{aligned}&\bar{\chi}^{A_{16},A_{15},A_{13},A_{23}}(\bar{x}_2(t_2)\bar{x}_3(t_3)\bar{x}_4(t_4)\bar{x}_5(t_5))\\ &:= \vartheta(A_{16}t_5)\cdot\vartheta\pi_q(A_{15}^q t_4 + A_{15}t_4^{q^2})\cdot\vartheta\pi_q(-A_{13}t_3)\cdot\vartheta(A_{23}t_2).\end{aligned}$$

Denote by $\psi^{A_{16},A_{15},A_{13},A_{23}}$ the lift of $\bar{\chi}^{A_{16},A_{15},A_{13},A_{23}}$ to T, then

$$\mathrm{Irr}(T)_1 := \{\psi \in \mathrm{Irr}(T) \mid X_6 \subseteq \ker\psi\} = \left\{\psi^{A_{16},A_{15},A_{13},A_{23}} \,\middle|\, A_{16},A_{23} \in \mathbb{F}_q, A_{13},A_{15} \in \mathbb{F}_{q^3}\right\}.$$

(2) If $X_6 \nsubseteq \ker\psi$, let $\psi^{A_{17}^*} := \mathrm{Ind}_N^T\lambda^{A_{17}^*,0,0}$. Then

$$\mathrm{Irr}(T)_2 := \{\psi \in \mathrm{Irr}(T) \mid X_6 \nsubseteq \ker\psi\} = \left\{\psi^{A_{17}^*} \,\middle|\, A_{17} \in \mathbb{F}_q^*\right\}.$$

Thus, $\mathrm{Irr}(T) = \mathrm{Irr}(T)_1\dot{\cup}\mathrm{Irr}(T)_2$, i.e. T has q^8 linear characters and $(q-1)$ irreducible characters of degree q^4.

By the commutator relations, we also determine the conjugacy classes of T and the character table of T.

4.11 Corollary. *Let $x := x(0,t_2,t_3,t_4,t_5,t_6) \in T$ be a representative of one conjugacy class of T, then the character table of T is the one in Table 3.*

Now we give the explicit constructions of the irreducible characters of U.

4.12 Proposition. *Let $A_{12},A_{13},A_{15} \in \mathbb{F}_{q^3}$, and $A_{23},A_{16},A_{17} \in \mathbb{F}_q$.*
Let $A_{12}^*,A_{13}^*,A_{15}^* \in \mathbb{F}_{q^3}^*$, $A_{23}^*,A_{16}^*,A_{17}^* \in \mathbb{F}_q^*$ and $T^{A_{13}^*}$ be a transversal of $A_{13}^*\mathbb{F}_q^+$ in $\mathbb{F}_{q^3}^+$.

Table 3: Character table of T

$\lvert^T x\rvert$	1	q	q	q	q
x	$x_6(t_6)$	$x_5(t_5^*)$	$x_4(t_4^*)x_5(t_5)$	$x_3(t_3^*)x_4(t_4)x_5(t_5)$	$x_2(t_2^*)x_3(t_3)$ $\cdot x_4(t_4)x_5(t_5)$
$\psi^{A_{16},A_{15},A_{13},A_{23}}$	1	$\vartheta(A_{16}t_5^*)$	$\vartheta(A_{16}t_5)$ $\cdot\vartheta\pi_q(A_{15}^q t_4^*)$ $\cdot\vartheta\pi_q(A_{15}t_4^{*q^2})$	$\vartheta(A_{16}t_5)$ $\cdot\vartheta\pi_q(A_{15}^q t_4)$ $\cdot\vartheta\pi_q(A_{15}t_4^{q^2})$ $\cdot\vartheta\pi_q(-A_{13}t_3^*)$	$\vartheta(A_{16}t_5)$ $\cdot\vartheta\pi_q(A_{15}^q t_4)$ $\cdot\vartheta\pi_q(A_{15}t_4^{q^2})$ $\cdot\vartheta\pi_q(-A_{13}t_3)$ $\cdot\vartheta(A_{23}t_2^*)$
$\psi^{A_{17}^*}$	$q^4\vartheta(A_{17}^*t_6)$	0	0	0	0

(1) Let $\bar{U} := X_3X_4X_5X_6\backslash U = \bar{X}_2\bar{X}_1$, $\bar{\chi}_{lin}^{A_{12},A_{23}} \in \mathrm{Irr}(\bar{U})$, $\bar{\chi}_{lin}^{A_{12},A_{23}}(\bar{x}_2(t_2)\bar{x}_1(t_1)) := \vartheta\pi_q(A_{12}t_1)\cdot\vartheta(A_{23}t_2)$, *and denote by* $\chi_{lin}^{A_{12},A_{23}}$ *the lift of* $\bar{\chi}_{lin}^{A_{12},A_{23}}$ *to* U, *then*

$$\mathfrak{F}_{lin} := \{\chi \in \mathrm{Irr}(U) \mid X_3X_4X_5X_6 \subseteq \ker\chi\} = \left\{\chi_{lin}^{A_{12},A_{23}} \,\middle|\, A_{12} \in \mathbb{F}_{q^3}, A_{23} \in \mathbb{F}_q\right\}.$$

(2) Let $\bar{A}_{12}^{A_{13}^*} \in T^{A_{13}^*}$, $\bar{U} := X_4X_5X_6\backslash U = \bar{X}_2\bar{X}_1\bar{X}_3$, $\bar{H} := \bar{X}_1\bar{X}_3$, $\bar{\chi}_{3,q}^{A_{13},A_{12}} \in \mathrm{Irr}(\bar{H})$ *with* $\bar{\chi}_{3,q}^{A_{13},A_{12}}(\bar{x}_1(t_1)\bar{x}_3(t_3)) := \vartheta\pi_q(A_{12}t_1 - A_{13}t_3)$. *Let* $\chi_{3,q}^{A_{13}^*,\bar{A}_{12}^{A_{13}^*}}$ *be the lift of* $\mathrm{Ind}_{\bar{H}}^{\bar{U}}\bar{\chi}_{3,q}^{A_{13}^*,\bar{A}_{12}^{A_{13}^*}}$ *to* U, *then*

$$\mathfrak{F}_3 := \{\chi \in \mathrm{Irr}(U) \mid X_4X_5X_6 \subseteq \ker\chi,\ X_3 \nsubseteq \ker\chi\} = \left\{\chi_{3,q}^{A_{13}^*,\bar{A}_{12}^{A_{13}^*}} \,\middle|\, A_{13}^* \in \mathbb{F}_{q^3}^*, \bar{A}_{12}^{A_{13}^*} \in T^{A_{13}^*}\right\}.$$

(3) Let $\bar{U} := X_5X_6\backslash U = \bar{X}_2\bar{X}_1\bar{X}_3\bar{X}_4$, $\bar{H} := \bar{X}_2\bar{X}_3\bar{X}_4$, $\bar{\chi}_{4,q^3}^{A_{15},A_{23},A_{13}} \in \mathrm{Irr}(\bar{H})$, *and*

$$\bar{\chi}_{4,q^3}^{A_{15},A_{23},A_{13}}(\bar{x}_2(t_2)\bar{x}_3(t_3)\bar{x}_4(t_4)) := \vartheta(A_{23}t_2)\cdot\vartheta\pi_q(-A_{13}t_3)\cdot\vartheta\pi_q(A_{15}^q t_4 + A_{15}t_4^{q^2}).$$

Denote by $\chi_{4,q^3}^{A_{15}^*,A_{23}}$ *the lift of* $\mathrm{Ind}_{\bar{H}}^{\bar{U}}\bar{\chi}_{4,q^3}^{A_{15}^*,A_{23},0}$ *to* U, *then*

$$\mathfrak{F}_4 := \{\chi \in \mathrm{Irr}(U) \mid X_5X_6 \subseteq \ker\chi,\ X_4 \nsubseteq \ker\chi\} = \left\{\chi_{4,q^3}^{A_{15}^*,A_{23}} \,\middle|\, A_{15}^* \in \mathbb{F}_{q^3}^*, A_{23} \in \mathbb{F}_q\right\}.$$

(4) Let $\bar{U} := X_6\backslash U = \bar{X}_2\bar{X}_1\bar{X}_3\bar{X}_4\bar{X}_5$, $\bar{H} := \bar{X}_2\bar{X}_3\bar{X}_4\bar{X}_5$, $\bar{\chi}_{5,q^3}^{A_{16},A_{23},A_{13},A_{15}} \in \mathrm{Irr}(\bar{H})$, *and*

$$\begin{aligned}&\bar{\chi}_{5,q^3}^{A_{16},A_{23},A_{13},A_{15}}(\bar{x}_2(t_2)\bar{x}_3(t_3)\bar{x}_4(t_4)\bar{x}_5(t_5))\\ :=&\vartheta(A_{23}t_2)\cdot\vartheta\pi_q(-A_{13}t_3)\cdot\vartheta\pi_q(A_{15}^q t_4 + A_{15}t_4^{q^2})\cdot\vartheta(A_{16}t_5).\end{aligned}$$

Denote by $\chi_{5,q^3}^{A_{16}^*,A_{23},A_{13}}$ *the lift of* $\mathrm{Ind}_{\bar{H}}^{\bar{U}}\bar{\chi}_{5,q^3}^{A_{16}^*,A_{23},A_{13},0}$ *to* U, *then*

$$\mathfrak{F}_5 := \{\chi \in \mathrm{Irr}(U) \mid X_6 \subseteq \ker\chi,\ X_5 \nsubseteq \ker\chi\} = \left\{\chi_{5,q^3}^{A_{16}^*,A_{23},A_{13}} \,\middle|\, A_{16}^* \in \mathbb{F}_q^*, A_{23} \in \mathbb{F}_q, A_{13} \in \mathbb{F}_{q^3}\right\}.$$

(5) Let $H := X_1X_4X_5X_6$, $\bar{H} := X_4X_5\backslash H \cong \bar{X}_1\bar{X}_6$, $\bar{\chi}_{6,q^4}^{A_{17},A_{12}} \in \mathrm{Irr}(\bar{H})$, *and*

$$\bar{\chi}_{6,q^4}^{A_{17},A_{12}}(\bar{x}_1(t_1)\bar{x}_6(t_6)) := \vartheta\pi_q(A_{12}t_1)\cdot\vartheta(A_{17}t_6).$$

Let $\tilde{\chi}_{6,q^4}^{A_{17},A_{12}}$ *denote the lift of* $\bar{\chi}_{6,q^4}^{A_{17},A_{12}}$ *from* $\bar{H}$ *to* H, *and* $\chi_{6,q^4}^{A_{17}^*,A_{12}} := \mathrm{Ind}_H^U\tilde{\chi}_{6,q^4}^{A_{17}^*,A_{12}}$, *then*

$$\mathfrak{F}_6 := \{\chi \in \mathrm{Irr}(U) \mid X_6 \nsubseteq \ker\chi\} = \left\{\chi_{6,q^4}^{A_{17}^*,A_{12}} \,\middle|\, A_{17}^* \in \mathbb{F}_q^*, A_{12} \in \mathbb{F}_{q^3}\right\}.$$

Hence $\mathrm{Irr}(U) = \mathfrak{F}_{lin} \dot\cup \mathfrak{F}_3 \dot\cup \mathfrak{F}_4 \dot\cup \mathfrak{F}_5 \dot\cup \mathfrak{F}_6$.

Proof. Let $\chi \in \mathrm{Irr}(U)$.

(1) *Family* $\mathfrak{F}_{lin}$*, where* $X_3X_4X_5X_6 \subseteq \ker\chi$.

Since $U' = X_3X_4X_5X_6$, all linear characters of U are precisely the lifts of the linear characters of the quotient group $X_3X_4X_5X_6\backslash U$.

(2) *Family* $\mathfrak{F}_3$*, where* $X_4X_5X_6 \subseteq \ker\chi$ *and* $X_3 \nsubseteq \ker\chi$.

Let $\bar U := X_4X_5X_6\backslash U = \bar X_2\bar X_1\bar X_3$, then the commutator relation in $\bar U$ is $[\bar x_2(t_2), \bar x_1(t_1)] = \bar x_3(t_2t_1)$. Let $\bar H := \bar X_1\bar X_3$, $\bar\chi_{3,q}^{A_{13},A_{12}} \in \mathrm{Irr}(\bar H)$ and

$$\bar\chi_{3,q}^{A_{13},A_{12}}(\bar x_1(t_1)\bar x_3(t_3)) := \vartheta\pi_q(A_{12}t_1)\cdot\vartheta\pi_q(-A_{13}t_3).$$

We note that $\bar X_2$ is a transversal of $\bar H$ in $\bar U$, and $Z(\bar U) = \bar X_3$. For all $s_2 \in \mathbb{F}_q$,

$$\begin{aligned}&\left(\bar\chi_{3,q}^{A_{13}^*,A_{12}}\right)^{\bar x_2(s_2)}(\bar x_1(t_1)\bar x_3(t_3)) = \bar\chi_{3,q}^{A_{13}^*,A_{12}}(\bar x_2(s_2)\cdot\bar x_1(t_1)\bar x_3(t_3)\cdot\bar x_2(s_2)^{-1})\\ =&\bar\chi_{3,q}^{A_{13}^*,A_{12}}(\bar x_1(t_1)\bar x_3(t_3+s_2t_1)) = \vartheta\pi_q(A_{12}t_1 - A_{13}^*(t_3+s_2t_1))\\ =&\vartheta\pi_q((A_{12}-s_2A_{13}^*)t_1 - A_{13}^*t_3) = \vartheta\pi_q((A_{12}-s_2A_{13}^*)t_1)\cdot\vartheta\pi_q(-A_{13}^*t_3),\end{aligned}$$

so $I_{\bar U}(\bar\chi_{3,q}^{A_{13}^*,A_{12}}) = \bar H$. By Clifford theory, $\tilde\chi_{3,q}^{A_{13}^*,A_{12}} := \mathrm{Ind}_{\bar H}^{\bar U}\bar\chi_{3,q}^{A_{13}^*,A_{12}} \in \mathrm{Irr}(\bar U)$ and

$$\mathrm{Res}_{\bar H}^{\bar U}\tilde\chi_{3,q}^{A_{13}^*,A_{12}} = \sum_{s_2\in\mathbb{F}_q}\left(\bar\chi_{3,q}^{A_{13}^*,A_{12}}\right)^{\bar x_2(s_2)} = \sum_{s_2\in\mathbb{F}_q}\bar\chi_{3,q}^{A_{13}^*,A_{12}-s_2A_{13}^*}.$$

By Clifford theory, there are $q^2(q^3-1)$ almost faithful irreducible characters of $\bar U$, i.e. $\{\tilde\chi_{3,q}^{A_{13}^*,\bar A_{12}^{A_{13}^*}} \mid A_{13}^* \in \mathbb{F}_{q^3}^*, \bar A_{12}^{A_{13}^*} \in T^{A_{13}^*}\}$. Let $\chi_{3,q}^{A_{13}^*,\bar A_{12}^{A_{13}^*}}$ be the lift of $\tilde\chi_{3,q}^{A_{13}^*,\bar A_{12}^{A_{13}^*}}$ to U. Thus,

$$\mathfrak{F}_3 = \{\chi\in\mathrm{Irr}(U) \mid X_4X_5X_6 \subseteq \ker\chi,\ X_3 \nsubseteq \ker\chi\} = \left\{\chi_{3,q}^{A_{13}^*,\bar A_{12}^{A_{13}^*}} \,\middle|\, A_{13}^* \in \mathbb{F}_{q^3}^*, \bar A_{12}^{A_{13}^*} \in T^{A_{13}^*}\right\}.$$

(3) The proof of the Family $\mathfrak{F}_4$ is an adaptation of the proof of the Family $\mathfrak{F}_3$.

(4) The proof of the Family $\mathfrak{F}_5$ is also an adaptation of the proof of the Family $\mathfrak{F}_3$.

(5) *Family* $\mathfrak{F}_6$*, where* $X_6 \nsubseteq \ker\chi$.

Let $T := X_2X_3X_4X_5X_6$, $N := X_4X_5X_6$, and $\chi \in \mathrm{Irr}(U)$ such that $X_6 \nsubseteq \ker(\chi)$, then $Z(T) = Z(U) = X_6$. Let $\psi \in \mathrm{Irr}(T)$ and $\langle\psi, \mathrm{Res}_T^U\chi\rangle_T > 0$, then $X_6 \nsubseteq \ker\psi$ by 4.4. Let $\lambda^{A_{17},A_{16},A_{15}} \in \mathrm{Irr}(N)$ and

$$\lambda^{A_{17},A_{16},A_{15}}(x_4(t_4)x_5(t_5)x_6(t_6)) := \vartheta(A_{17}t_6)\cdot\vartheta(A_{16}t_5)\cdot\vartheta\pi_q(A_{15}^qt_4 + A_{15}t_4^{q^2}),$$

then by 4.10, $\{\psi\in\mathrm{Irr}(T) \mid X_6 \nsubseteq \ker\psi\} = \{\mathrm{Ind}_N^T\lambda^{A_{17}^*,0,0} \mid A_{17}^* \in \mathbb{F}_q^*\} = \{\psi^{A_{17}^*} \mid A_{17}^* \in \mathbb{F}_q^*\}$. By (5) of 4.6, $I_U(\psi^{A_{17}^*}) = U$, so $\mathrm{Res}_T^U\chi = z^*\psi^{A_{17}^*}$ for some $z^* \in \mathbb{N}^*$. Thus

$$\begin{aligned}\mathfrak{F}_6 =&\{\chi\in\mathrm{Irr}(U) \mid X_6 \nsubseteq \ker\chi\} = \bigcup_{\substack{\psi\in\mathrm{Irr}(T)\\ X_6\nsubseteq\ker\psi}}\{\chi\in\mathrm{Irr}(U) \mid \langle\chi, \mathrm{Ind}_T^U\psi\rangle_U > 0\}\\ =&\bigcup_{A_{17}^*\in\mathbb{F}_q^*}\{\chi\in\mathrm{Irr}(U) \mid \langle\chi, \mathrm{Ind}_T^U\psi^{A_{17}^*}\rangle_U > 0\}\\ =&\bigcup_{A_{17}^*\in\mathbb{F}_q^*}\{\chi\in\mathrm{Irr}(U) \mid \langle\chi, \mathrm{Ind}_N^U\lambda^{A_{17}^*,0,0}\rangle_U > 0\}.\end{aligned}$$

Let $H := X_1X_4X_5X_6$, then $H' = X_5$ and $Z(H) = X_4X_5 \trianglelefteq H$. Let $\tilde{\chi}^{A_{17},A_{15},A_{12}} \in \mathrm{Irr}(H)$ as in (1) of 4.8. For all $x_4(t_4)x_5(t_5)x_6(t_6) \in N$,

$$\begin{aligned}&\left(\mathrm{Res}_N^H \tilde{\chi}^{A_{17}^*,0,A_{12}}\right)(x_4(t_4)x_5(t_5)x_6(t_6)) = \tilde{\chi}^{A_{17}^*,0,A_{12}}(x_4(t_4)x_5(t_5)x_6(t_6))\\ =&\bar{\chi}(\bar{x}_4(t_4)\bar{x}_6(t_6)) = \vartheta(A_{17}^* t_6) = \lambda^{A_{17}^*,0,0}(x_4(t_4)x_5(t_5)x_6(t_6)).\end{aligned}$$

Thus $\mathrm{Res}_N^H \tilde{\chi}^{A_{17}^*,0,A_{12}} = \lambda^{A_{17}^*,0,0}$ for all $A_{12} \in \mathbb{F}_{q^3}$. By (4) of 4.6 we get $I_H(\lambda^{A_{17}^*,0,0}) = H$, thus $\mathrm{Ind}_N^H \lambda^{A_{17}^*,0,0} = \sum_{A_{12}\in\mathbb{F}_{q^3}} \tilde{\chi}^{A_{17}^*,0,A_{12}}$. By (2) of 4.6 $I_U(\lambda^{A_{17}^*,0,0}) = H$. Then by Clifford's Theorem, $\mathrm{Ind}_H^U \tilde{\chi}^{A_{17}^*,0,A_{12}} \in \mathrm{Irr}(U)$ for all $A_{17}^* \in \mathbb{F}_q^*$. Thus

$$\begin{aligned}\mathfrak{F}_6 &= \bigcup_{A_{17}^*\in\mathbb{F}_q^*} \{\chi \in \mathrm{Irr}(U) \mid \langle \chi, \mathrm{Ind}_H^U \mathrm{Ind}_N^H \lambda^{A_{17}^*,0,0}\rangle_U > 0\}\\ &= \bigcup_{\substack{A_{17}^*\in\mathbb{F}_q^*\\ A_{12}\in\mathbb{F}_{q^3}}} \{\chi \in \mathrm{Irr}(U) \mid \langle \chi, \mathrm{Ind}_H^U \tilde{\chi}^{A_{17}^*,0,A_{12}}\rangle_U > 0\}\\ &= \left\{\mathrm{Ind}_H^U \tilde{\chi}^{A_{17}^*,0,A_{12}} \,\middle|\, A_{17}^* \in \mathbb{F}_q^*, A_{12} \in \mathbb{F}_{q^3}\right\}.\end{aligned}$$

For $A_{17}^* \in \mathbb{F}_q^*$ and $A_{12} \in \mathbb{F}_{q^3}$, $X_4X_5 \subseteq \ker(\tilde{\chi}^{A_{17}^*,0,A_{12}})$ and $X_4X_5 \trianglelefteq H$, so $\tilde{\chi}^{A_{17}^*,0,A_{12}}$ is the lift to H of some irreducible character of $\bar{H} := X_4X_5\backslash H \cong \bar{X}_1\bar{X}_6$. Let $\lambda^{A_{17},A_{16},A_{15}} \in \mathrm{Irr}(N)$, $\bar{\chi}_{6,q^4}^{A_{17}^*,A_{12}} \in \mathrm{Irr}(\bar{H})$, and $\bar{\chi}_{6,q^4}^{A_{17}^*,A_{12}}(\bar{x}_1(t_1)\bar{x}_6(t_6)) := \vartheta\pi_q(A_{12}t_1)\cdot\vartheta(A_{17}^* t_6)$. Let $\tilde{\chi}_{6,q^4}^{A_{17}^*,A_{12}}$ denote the lift of $\bar{\chi}_{6,q^4}^{A_{17}^*,A_{12}}$ from $\bar{H}$ to H, then $\tilde{\chi}_{6,q^4}^{A_{17}^*,A_{12}} = \tilde{\chi}^{A_{17}^*,0,A_{12}}$.

Let $\chi_{6,q^4}^{A_{17}^*,A_{12}} := \mathrm{Ind}_H^U \tilde{\chi}_{6,q^4}^{A_{17}^*,A_{12}}$, then

$$\mathfrak{F}_6 = \left\{\mathrm{Ind}_H^U \tilde{\chi}^{A_{17}^*,0,A_{12}} \,\middle|\, A_{17}^* \in \mathbb{F}_q^*, A_{12} \in \mathbb{F}_{q^3}\right\} = \left\{\chi_{6,q^4}^{A_{17}^*,A_{12}} \,\middle|\, A_{17}^* \in \mathbb{F}_q^*, A_{12} \in \mathbb{F}_{q^3}\right\}.$$

□

4.13 Remark. *From the proof (5) of Propositon 4.12, we obtain that*

$$\begin{aligned}&\mathrm{Ind}_T^U \psi^{A_{17}^*} = \mathrm{Ind}_N^U \lambda^{A_{17}^*,0,0} = \mathrm{Ind}_H^U\left(\sum_{A_{12}\in\mathbb{F}_q} \tilde{\chi}^{A_{17}^*,0,A_{12}}\right)\\ =&\mathrm{Ind}_H^U\left(\sum_{A_{12}\in\mathbb{F}_q} \tilde{\chi}_{6,q^4}^{A_{17}^*,A_{12}}\right) = \sum_{A_{12}\in\mathbb{F}_q} \mathrm{Ind}_H^U \tilde{\chi}_{6,q^4}^{A_{17}^*,A_{12}} = \sum_{A_{12}\in\mathbb{F}_q} \chi_{6,q^4}^{A_{17}^*,A_{12}}\end{aligned}$$

5 Character table

In this section, we determine the character table of ${}^3D_4^{syl}(q^3)$. We use the notations as in Section 4.

5.1 Proposition. *The character table of ${}^3D_4^{syl}(q^3)$ is the one in Table 4.*

Table 4: Character table of ${}^3D_4^{syl}(q^3)$

	I_8	$x_1(t_1^*)$ $\cdot x_6(t_6)$	$x_1(t_1^*)$ $\cdot x_3(\bar{t}_3^*)$	$x_2(t_2^*)$ $\cdot x_4(t_4)$ $\cdot x_5(t_5)$	$x_2(t_2^*)$ $\cdot x_1(t_1^*)$	$x_3(t_3^*)$ $\cdot x_5(t_5)$	$x_4(t_4^*)$	$x_5(t_5^*)$	$x_6(t_6^*)$
$\chi_{lin}^{0,0}$	1	1	1	1	1	1	1	1	1
$\chi_{lin}^{A_{12}^*,0}$	1	$\vartheta\pi_q(A_{12}^*t_1^*)$	$\vartheta\pi_q(A_{12}^*t_1^*)$	1	$\vartheta\pi_q(A_{12}^*t_1^*)$	1	1	1	1
χ_{lin}^{0,A_{23}^*}	1	1	1	$\vartheta(A_{23}^*t_2^*)$	$\vartheta(A_{23}^*t_2^*)$	1	1	1	1
$\chi_{lin}^{A_{12}^*,A_{23}^*}$	1	$\vartheta\pi_q(A_{12}^*t_1^*)$	$\vartheta\pi_q(A_{12}^*t_1^*)$	$\vartheta(A_{23}^*t_2^*)$	$\vartheta\pi_q(A_{12}^*t_1^*)$ $\cdot\vartheta(A_{23}^*t_2^*)$	1	1	1	1
$\chi_{3,q}^{A_{13}^*,\bar{A}_{12}^{A_{13}^*}}$	q	$\divideontimes$	$\divideontimes$	0	0	$\vartheta\pi_q(-A_{13}^*t_3^*)$ $\cdot q$	q	q	q
$\chi_{4,q^3}^{A_{15}^*,A_{23}}$	q^3	0	0	$\divideontimes$	0	0	$\vartheta\pi_q(A_{15}^{*q}t_4^*)$ $\cdot\vartheta\pi_q(A_{15}^*t_4^{*q^2})$ $\cdot q^3$	q^3	q^3
$\chi_{5,q^3}^{A_{16}^*,A_{23},A_{13}}$	q^3	0	0	$\divideontimes$	0	$\divideontimes$	0	$\vartheta(A_{16}^*t_5^*)$ $\cdot q^3$	q^3
$\chi_{6,q^4}^{A_{17}^*,A_{12}}$	q^4	$\divideontimes$	0	0	0	0	0	0	$\vartheta(A_{17}^*t_6^*)$ $\cdot q^4$

where $\bar{A}_{12}^{A_{13}^*} \in T^{A_{13}^*}$, $0 \neq \bar{t}_3^* \in T^{t_1^*}$, and

$$\begin{aligned}\chi_{6,q^4}^{A_{17}^*,A_{12}}(x_1(t_1^*)x_6(t_6)) &= \sum_{r_3\in\mathbb{F}_{q^3}} \vartheta\pi_q\left(A_{12}t_1^* + A_{17}^*\phi_0(-t_1^* r_3^{q^2+q}) + A_{17}^* t_6\right)\\ &= \vartheta\pi_q(A_{12}t_1^* + A_{17}^* t_6)\cdot \sum_{r_3\in\mathbb{F}_{q^3}} \vartheta\left(A_{17}^*\phi_0(-t_1^* r_3^{q^2+q})\right),\\ \chi_{5,q^3}^{A_{16}^*,A_{23},A_{13}}(x_3(t_3^*)x_5(t_5)) &= \sum_{r_1\in\mathbb{F}_{q^3}} \vartheta\pi_q\left(-A_{13}t_3^* + A_{16}^*\phi_0(t_3^* r_1^{q^2+q}) + A_{16}^* t_5\right)\\ &= \vartheta\pi_q(-A_{13}t_3^* + A_{16}^* t_5)\cdot \sum_{r_1\in\mathbb{F}_{q^3}} \vartheta\left(A_{16}^*\phi_0(t_3^* r_1^{q^2+q})\right),\end{aligned}$$

$$\begin{aligned}&\chi_{5,q^3}^{A_{16}^*,A_{23},A_{13}}(x_2(t_2^*)x_4(t_4)x_5(t_5))\\ =&\sum_{r_1\in\mathbb{F}_{q^3}} \vartheta\pi_q(A_{23}t_2^* + A_{13}r_1 t_2^* + A_{16}^*(\phi_0(r_1 t_4^q) - t_2^* r_1^{q^2+q+1}) + A_{16}^* t_5)\\ =&\vartheta(A_{23}t_2^* + A_{16}^* t_5)\cdot \sum_{r_1\in\mathbb{F}_{q^3}} \vartheta\pi_q\left(A_{13}r_1 t_2^* + A_{16}^*(\phi_0(r_1 t_4^q) - t_2^* r_1^{q^2+q+1})\right),\end{aligned}$$

$$\begin{aligned}&\chi_{4,q^3}^{A_{15}^*,A_{23}}(x_2(t_2^*)x_4(t_4)x_5(t_5))\\ =&\sum_{r_1\in\mathbb{F}_{q^3}} \vartheta\pi_q(A_{23}t_2^* + {A_{15}^*}^q(t_4 - t_2^* r_1^{q+1}) + A_{15}^*(t_4^{q^2} - t_2^* r_1^{q^2+1}))\\ =&\vartheta(A_{23}t_2^*)\cdot \sum_{r_1\in\mathbb{F}_{q^3}} \vartheta\pi_q\left({A_{15}^*}^q(t_4 - t_2^* r_1^{q+1}) + A_{15}^*(t_4^{q^2} - t_2^* r_1^{q^2+1})\right),\end{aligned}$$

$$\begin{aligned}\chi_{3,q}^{A_{13}^*,\bar{A}_{12}^{A_{13}^*}}(x_1(t_1^*)x_6(t_6)) &= \sum_{r_2\in\mathbb{F}_q} \vartheta\pi_q\left((\bar{A}_{12}^{A_{13}^*} - A_{13}^* r_2)t_1^*\right),\\ \chi_{3,q}^{A_{13}^*,\bar{A}_{12}^{A_{13}^*}}(x_1(t_1^*)x_3(\bar{t}_3^*)) &= \sum_{r_2\in\mathbb{F}_q} \vartheta\pi_q\left((\bar{A}_{12}^{A_{13}^*} - A_{13}^* r_2)t_1^* - \bar{t}_3^* A_{13}^*\right).\end{aligned}$$

Proof. Let $x := x(t_1,t_2,t_3,t_4,t_5,t_6) \in U = {}^3D_4^{syl}(q^3)$.

(1) *Calculate the values of* $\chi_{6,q^4}^{A_{17}^*,A_{12}}$ *for all* $A_{17}^* \in \mathbb{F}_q^*$ *and* $A_{12} \in \mathbb{F}_{q^3}$.

We use the notations of (5) of Proposition 4.12, then

$$\begin{aligned}\chi_{6,q^4}^{A_{17}^*,A_{12}}(x) = \left(\mathrm{Ind}_H^U \tilde{\chi}_{6,q^4}^{A_{17}^*,A_{12}}\right)(x) &= \frac{1}{|H|}\sum_{\substack{g\in U\\ g\cdot x\cdot g^{-1}\in H}} \tilde{\chi}_{6,q^4}^{A_{17}^*,A_{12}}(g\cdot x\cdot g^{-1})\\ =&\frac{1}{|H|}\sum_{\substack{g\in U\\ g\cdot x\cdot g^{-1}\in H}} \bar{\chi}_{6,q^4}^{A_{17}^*,A_{12}}(X_4X_5\backslash(g\cdot x\cdot g^{-1})).\end{aligned}$$

Thus,

$$\begin{aligned}&\chi_{6,q^4}^{A_{17}^*,A_{12}}(x_1(t_1^*)x_3(\bar{t}_3^*)) = \chi_{6,q^4}^{A_{17}^*,A_{12}}(x_2(t_2^*)x_4(t_4)x_5(t_5))\\ =&\chi_{6,q^4}^{A_{17}^*,A_{12}}(x_2(t_2^*)x_1(t_1^*)) = \chi_{6,q^4}^{A_{17}^*,A_{12}}(x_3(t_3^*)x_5(t_5)) \overset{gxg^{-1}\notin H}{=} 0.\end{aligned}$$

Then

$$\begin{aligned}
&\chi_{6,q^4}^{A_{17}^*,A_{12}}\left(x_4(t_4)x_5(t_5)x_6(t_6)\right)\\
=&\frac{1}{|H|}\sum_{\substack{g:=x(r_1,r_2,r_3,r_4,r_5,r_6)\in U\\ g\cdot x_4(t_4)x_5(t_5)x_6(t_6)\cdot g^{-1}\in H}}\tilde{\chi}_{6,q^4}^{A_{17}^*,A_{12}}(g\cdot x_4(t_4)x_5(t_5)x_6(t_6)\cdot g^{-1})\\
=&\frac{1}{|H|}\sum_{\substack{r_1,r_3,r_4\in\mathbb{F}_{q^3}\\ r_2,r_5,r_6\in\mathbb{F}_q}}\tilde{\chi}_{6,q^4}^{A_{17}^*,A_{12}}(x_4(t_4)x_5(t_5+\phi_0\left(r_1t_4^q\right))\\
&\qquad\qquad\cdot x_6(t_6+r_2t_5+\phi_0(r_1r_2t_4^q)+\phi_0(r_3t_4^q)))\\
=&\frac{1}{|H|}\sum_{\substack{r_1,r_3,r_4\in\mathbb{F}_{q^3}\\ r_2,r_5,r_6\in\mathbb{F}_q}}\bar{\chi}_{6,q^4}^{A_{17}^*,A_{12}}(\bar{x}_6\left(t_6+r_2t_5+\phi_0(r_1r_2t_4^q)+\phi_0(r_3t_4^q)\right))\\
=&\frac{1}{q^3}\sum_{\substack{r_1,r_3\in\mathbb{F}_{q^3}\\ r_2\in\mathbb{F}_q}}\bar{\chi}_{6,q^4}^{A_{17}^*,A_{12}}(\bar{x}_6\left(t_6+r_2t_5+\phi_0(r_1r_2t_4^q)+\phi_0(r_3t_4^q)\right)),
\end{aligned}$$

so,

$$\chi_{6,q^4}^{A_{17}^*,A_{12}}(I_8)=q^4,\ \chi_{6,q^4}^{A_{17}^*,A_{12}}(x_4(t_4^*))=\chi_{6,q^4}^{A_{17}^*,A_{12}}(x_5(t_5^*))=0,\ \chi_{6,q^4}^{A_{17}^*,A_{12}}(x_6(t_6^*))=q^4\cdot\vartheta(A_{17}^*t_6^*).$$

Then

$$\begin{aligned}
\chi_{6,q^4}^{A_{17}^*,A_{12}}\left(x_1(t_1^*)x_6(t_6)\right)=&\frac{1}{|H|}\sum_{\substack{g\in U\\ g\cdot x_1(t_1^*)x_6(t_6)\cdot g^{-1}\in H}}\tilde{\chi}_{6,q^4}^{A_{17}^*,A_{12}}(g\cdot x_1(t_1^*)x_6(t_6)\cdot g^{-1})\\
\overset{g:=x(r_1,r_2,r_3,r_4,r_5,r_6)}{=}&\frac{1}{|H|}\sum_{\substack{r_2=0\\ r_1,r_3,r_4\in\mathbb{F}_{q^3}\\ r_5,r_6\in\mathbb{F}_q}}\bar{\chi}_{6,q^4}^{A_{17}^*,A_{12}}(\bar{x}_1(t_1^*)\bar{x}_6(t_6+\phi_0\left(-t_1^*r_3^{q^2+q}\right)))\\
=&\sum_{r_3\in\mathbb{F}_{q^3}}\bar{\chi}_{6,q^4}^{A_{17}^*,A_{12}}(\bar{x}_1(t_1^*)\bar{x}_6(t_6+\phi_0\left(-t_1^*r_3^{q^2+q}\right)))\\
=&\vartheta\pi_q(A_{12}t_1^*+A_{17}^*t_6)\cdot\sum_{r_3\in\mathbb{F}_{q^3}}\vartheta\left(A_{17}^*\phi_0(-t_1^*r_3^{q^2+q})\right).
\end{aligned}$$

Thus all the values of $\chi_{6,q^4}^{A_{17}^*,A_{12}}$ are obtained.

(2) *Calculate the values of* $\chi_{3,q}^{A_{13}^*,\bar{A}_{12}^{A_{13}^*}}$ *for all* $A_{13}^*\in\mathbb{F}_{q^3}^*$ *and* $\bar{A}_{12}^{A_{13}^*}\in T^{A_{13}^*}$.

We use the notations of (2) of Proposition 4.12. For $x=x(t_1,t_2^*,t_3,t_4,t_5,t_6)\in U$ with $0\neq t_2^*\in\mathbb{F}_q^*$, we have

$$\chi_{3,q}^{A_{13}^*,\bar{A}_{12}^{A_{13}^*}}(x)=\left(\mathrm{Ind}_{\bar{H}}^{\bar{U}}\bar{\chi}_{3,q}^{A_{13}^*,\bar{A}_{12}^{A_{13}^*}}\right)(\bar{x})=\frac{1}{|\bar{H}|}\sum_{\substack{\bar{u}\in\bar{U}\\ \bar{u}\cdot\bar{x}\cdot\bar{u}^{-1}\in\bar{H}}}\bar{\chi}_{3,q}^{A_{13}^*,\bar{A}_{12}^{A_{13}^*}}(\bar{u}\cdot\bar{x}\cdot\bar{u}^{-1})\overset{\bar{u}\cdot\bar{x}\cdot\bar{u}^{-1}\notin\bar{H}}{=}0.$$

If $x = x(t_1, 0, t_3, t_4, t_5, t_6) \in U$, we have

$$\begin{aligned}
&\chi_{3,q}^{A_{13}^*,\bar{A}_{12}^{A_{13}^*}}\left(x(t_1,0,t_3,t_4,t_5,t_6)\right) = \left(\mathrm{Ind}_{\bar{H}}^{\bar{U}}\bar{\chi}_{3,q}^{A_{13}^*,\bar{A}_{12}^{A_{13}^*}}\right)\left(\bar{x}_1(t_1)\bar{x}_3(t_3)\right)\\
=&\frac{1}{|\bar{H}|}\sum_{\substack{\bar{u}:=\bar{x}_2(r_2)\bar{x}_1(r_1)\bar{x}_3(r_3)\in\bar{U}\\ \bar{u}\cdot\bar{x}_1(t_1)\bar{x}_3(t_3)\cdot\bar{u}^{-1}\in\bar{H}}}\bar{\chi}_{3,q}^{A_{13}^*,\bar{A}_{12}^{A_{13}^*}}(\bar{u}\cdot\bar{x}_1(t_1)\bar{x}_3(t_3)\cdot\bar{u}^{-1})\\
=&\frac{1}{|\bar{H}|}\sum_{\substack{r_1,r_3\in\mathbb{F}_{q^3}\\ r_2\in\mathbb{F}_q}}\bar{\chi}_{3,q}^{A_{13}^*,\bar{A}_{12}^{A_{13}^*}}(\bar{x}_1(t_1)\bar{x}_3(t_3+r_2t_1))\\
=&\sum_{r_2\in\mathbb{F}_q}\bar{\chi}_{3,q}^{A_{13}^*,\bar{A}_{12}^{A_{13}^*}}(\bar{x}_1(t_1)\bar{x}_3(t_3+r_2t_1)) = \sum_{r_2\in\mathbb{F}_q}\vartheta\pi_q\left(\bar{A}_{12}^{A_{13}^*}t_1 - A_{13}^*(t_3+r_2t_1)\right).
\end{aligned}$$

Thus we get all the values of $\chi_{3,q}^{A_{13}^*,\bar{A}_{12}^{A_{13}^*}}$.

Similarly, we calculate all of the other values of the character table. □

In the paper, p is an odd prime. By Propositions 3.11, 4.12 and 5.1, we obtain the number of the conjugacy classes, and determine the numbers of the complex irreducible characters of degree q^c with $c \in \mathbb{N}$. T. Le [Le13] counted the number of irreducible characters for all primes p.

5.2 Corollary. *Let* $\#\mathrm{Irr}$ *be the number of irreducible characters of* ${}^3D_4^{syl}(q^3)$*, and* $\#\mathrm{Irr}_c$ *be the number of irreducible characters of* ${}^3D_4^{syl}(q^3)$ *of dimension* q^c *with* $c \in \mathbb{N}$*, then*

$$\begin{aligned}
\#\mathrm{Irr}_4 &= q^4 - q^3 &&= (q-1)^4 + 3(q-1)^3 + 3(q-1)^2 + (q-1),\\
\#\mathrm{Irr}_3 &= q^5 - q &&= (q-1)^5 + 5(q-1)^4 + 10(q-1)^3 + 10(q-1)^2 + 4(q-1),\\
\#\mathrm{Irr}_1 &= q^5 - q^2 &&= (q-1)^5 + 5(q-1)^4 + 10(q-1)^3 + 9(q-1)^2 + 3(q-1),\\
\#\mathrm{Irr}_0 &= q^4 &&= (q-1)^4 + 4(q-1)^3 + 6(q-1)^2 + 4(q-1) + 1,
\end{aligned}$$

and

$$\begin{aligned}
\#\mathrm{Irr} =&\#\{\textit{Irreducible Characters of } {}^3D_4^{syl}(q^3)\}\\
=&\#\{\textit{Conjugacy Classes of } {}^3D_4^{syl}(q^3)\}\\
=&2q^5 + 2q^4 - q^3 - q^2 - q\\
=&2(q-1)^5 + 12(q-1)^4 + 27(q-1)^3 + 28(q-1)^2 + 12(q-1) + 1.
\end{aligned}$$

5.3 Remark. *We also consider the analogue of Higman's conjecture, Lehrer's conjecture and Isaacs' conjecture of* $U_n(q)$ *for* ${}^3D_4^{syl}(q^3)$*. By 5.2, the conjectures are true for* ${}^3D_4^{syl}(q^3)$*.*

Acknowledgements

I would like to thank Richard Dipper, Mathias Werth and Yichuan Yang for the helpful discussions and valuable suggestions.

References

[Car72] Roger W. Carter. *Simple groups of Lie type*. John Wiley & Sons, New York, 1972.

[CR81] Charles W. Curtis and Irving Reiner. *Methods of representation theory: with applications to finite groups and orders*, volume I. John Wiley & Sons, Inc., New York, 1981.

[GLM15] Simon M. Goodwin, Tung Le, and Kay Magaard. The generic character table of a Sylow p-subgroup of a finite Chevalley group of type D_4. *preprint arXiv:1508.06937*, 2015.

[GLMP16] Simon M. Goodwin, Tung Le, Kay Magaard, and Alessandro Paolini. Constructing characters of Sylow p-subgroups of finite Chevalley groups. *J. Algebra*, 468:395–439, 2016.

[GMR14] Simon M. Goodwin, Peter Mosch, and Gerhard Röhrle. Calculating conjugacy classes in Sylow p-subgroups of finite Chevalley groups of rank six and seven. *LMS J. Comput. Math.*, 17(1):109–122, 2014.

[GMR16] Simon M. Goodwin, Peter Mosch, and Gerhard Röhrle. On the coadjoint orbits of maximal unipotent subgroups of reductive groups. *Transformation Groups*, 21(2):399–426, 2016.

[GR09] Simon M. Goodwin and Gerhard Röhrle. Calculating conjugacy classes in Sylow p-subgroups of finite Chevalley groups. *J. Algebra*, 321(11):3321–3334, 2009.

[Hig60] Graham Higman. Enumerating p-groups. I. Inequalities. *Proc. London Math. Soc. (3)*, 10:24–30, 1960.

[HLM11] Frank Himstedt, Tung Le, and Kay Magaard. Characters of the Sylow p-subgroups of the Chevalley groups $D_4(p^n)$. *J. Algebra*, 332:414–427, 2011.

[HLM16] Frank Himstedt, Tung Le, and Kay Magaard. On the characters of the Sylow p-subgroups of untwisted Chevalley groups $Y_n(p^a)$. *LMS J. Comput. Math.*, 19(2):303–359, 2016.

[Hum78] James E. Humphreys. *Introduction to Lie algebras and representation theory*, volume 9 of *Graduate Texts in Mathematics*. Springer-Verlag, New York-Berlin, 1978. Second printing, revised.

[Isa95] I. M. Isaacs. Characters of groups associated with finite algebras. *J. Algebra*, 177(3):708–730, 1995.

[Isa07] I. M. Isaacs. Counting characters of upper triangular groups. *J. Algebra*, 315(2):698–719, 2007.

[Jed13] Markus Jedlitschky. *Decomposing André-Neto supercharacters of Sylow p-subgroups of Lie type D*. PhD thesis, Universität Stuttgart, 2013.

[Le13] Tung Le. Irreducible characters of Sylow p-subgroups of the Steinberg triality groups ${}^3D_4(p^{3m})$. preprint arXiv:1309.1583, 2013.

[Leh74] G. I. Lehrer. Discrete series and the unipotent subgroup. *Compositio Math.*, 28:9–19, 1974.

[PS15] Igor Pak and Andrew Soffer. On Higman's $k(U_n(q))$ conjecture. preprint arXiv: 1507.00411v1, 2015.

[Sun16] Yujiao Sun. A supercharacter theory for Sylow p-subgroups ${}^3D_4^{syl}(q^3)$ of the Steinberg triality groups. preprint arXiv:1611.09287, 2016.

[VLA03] Antonio Vera-López and J. M. Arregi. Conjugacy classes in unitriangular matrices. *Linear Algebra Appl.*, 370:85–124, 2003.